

## 29:194 Homework #3

Due at the beginning of class, Thursday, September 18, 2008.

1. Show that the curvature drift

$$\mathbf{V}_c = \frac{v_{\parallel}^2}{\omega_c B} \frac{\mathbf{R}_c \times \mathbf{B}}{R_c^2}$$

can be written as

$$\mathbf{V}_c = \frac{v_{\parallel}^2}{\omega_c} \hat{\mathbf{b}} \times (\hat{\mathbf{b}} \cdot \nabla) \hat{\mathbf{b}}$$

You may take the magnetic field to be purely azimuthal  $\mathbf{B} = B \hat{\phi}$ .

HINT: Convert from cylindrical to Cartesian coordinates.

2. A 200 keV deuteron in a large mirror fusion device has a pitch angle of  $45^\circ$  at the midplane of the machine, where the magnetic field  $B = 0.7$  T. Compute its Larmor radius.
3. The equation for a dipole magnetic field in spherical coordinates is given by

$$\mathbf{B} = \frac{\mu_0 M}{4\pi} \frac{1}{r^3} (2 \cos \theta \hat{\mathbf{r}} + \sin \theta \hat{\boldsymbol{\theta}})$$

where  $M$  is the magnetic moment.

- (a) Show that the equation for a magnetic field line is  $r = R \sin^2 \theta$ , where  $R$  is the radius of the magnetic field line at the equator ( $\theta = \pi/2$ ).
- (b) Show that the curvature of the magnetic field line at the equator is  $R_C = R/3$ .
- (c) Compute the curvature drift of a particle with charge  $q$  and parallel velocity  $v_{\parallel}$  at a radial distance  $R$  at the equator.
- (d) Compute the  $\nabla B$  drift of a particle with charge  $q$  and perpendicular velocity  $v_{\perp}$  at a radial distance  $R$  at the equator.
- (e) Compare the equations for the curvature drift and the  $\nabla B$  drift at the equator.
4. A particle is trapped in a magnetic mirror field given by

$$B_z = B_0 \left[ 1 + \left( \frac{z}{L} \right)^2 \right]$$

and has a total kinetic energy  $w = mv^2/2$  and pitch angle  $\alpha$  at  $z = 0$ . Find the oscillation frequency in terms of  $L$ ,  $w$ , and  $\alpha$ .