

# PHYS:7729 Homework #4

Reading: Required: Read GB Chapter 9, Sections 9.1–9.3 (p.319–349)  
Optional: Read BS Chapter 7, Sections 7.1–7.3 (p.252–268)

Due at 5:00pm, Friday, March 26, 2021.

1. A plasma has a “spherical shell” distribution function given by

$$f_0(\mathbf{v}) = \frac{n_0}{4\pi C^2} \delta(|\mathbf{v}| - C)$$

where  $C$  is a constant.

- (a) Using the Fourier analysis approach, show that the dispersion relation for electrostatic waves in this plasma is  $\omega^2 = \omega_p^2 + k^2 C^2$ .  
(b) What is the region of validity of this dispersion relation?

2. Show that the Laplace transform of  $f(t) = \cosh(at)$  is given by

$$\tilde{f}(p) = \frac{p}{p^2 - a^2}.$$

3. Use the Residue Theorem to evaluate the inverse Laplace transform of

$$\tilde{f}(p) = \frac{1}{p^2 - a^2}.$$

4. Solution of Navier-Stokes Equations:

The Navier-Stokes Equations for the viscous evolution of a hydrodynamic fluid are given by:

$$\begin{aligned}\frac{\partial \rho}{\partial t} + \mathbf{U} \cdot \nabla \rho &= -\rho \nabla \cdot \mathbf{U} \\ \rho \left( \frac{\partial \mathbf{U}}{\partial t} + \mathbf{U} \cdot \nabla \mathbf{U} \right) &= -\nabla p + \nu \rho \nabla^2 \mathbf{U} \\ \frac{\partial p}{\partial t} + \mathbf{U} \cdot \nabla p &= -\gamma p \nabla \cdot \mathbf{U}\end{aligned}$$

where  $\nu$  is the coefficient of kinematic viscosity. Assume a wave vector of the form  $\mathbf{k} = k\hat{\mathbf{z}}$ . The initial conditions for a sound wave in this system at  $t = 0$  are  $\mathbf{U}(\mathbf{x}, 0) = \bar{U} \cos(kz)\hat{\mathbf{z}}$  and  $\mathbf{U}'(\mathbf{x}, 0) = -\bar{U}\omega_0 \sin(kz)\hat{\mathbf{z}}$ . Use the Laplace-Fourier transform method (Fourier transform in space, Laplace transform in time) to solve for the velocity  $\mathbf{U}(\mathbf{x}, t)$ . Note that the  $z$ -component of the velocity  $U_z$  is the only non-trivial part of the solution.

HINT: This is similar to a linear dispersion relation problem, so your first step is to linearize the Navier-Stokes equations.

- (a) Fourier transform the linearized equations and find the differential equation for  $U_z(\mathbf{k}, t)$  in terms of time derivatives. Use the definition of the sound speed  $c_s^2 = \gamma p_0 / \rho_0$  to simplify the equation.
- (b) Solve for the Laplace transform  $\tilde{U}_z(\mathbf{k}, p)$ .
- (c) Perform the inverse Laplace transform to find a solution for  $U_z(\mathbf{k}, t)$ . You may wish to define  $\omega^2 = k^2 c_s^2 - \nu^2 k^4 / 4$  to simplify notation.
- (d) Fourier transform the initial conditions and apply them to the answer above so that you may obtain the final solution  $U_z(\mathbf{x}, t)$ .
- (e) Determine the evolution of the magnitude of the velocity  $|U_z(\mathbf{x}, t)|$  for  $\nu^2 k^2 < 4c_s^2$ .
- (f) In the weak damping limit  $\nu^2 k^2 \ll 4c_s^2$ , what are the effective real frequency of oscillation (include the small, first order correction) and damping rate?
- (g) Qualitatively sketch the solution  $U_z(z = 0, t)$  in the case that  $\nu^2 k^2 < 4c_s^2$ .
- (h) Qualitatively sketch  $U_z(z = 0, t)$  for the cases  $\nu^2 k^2 = 4c_s^2$  and  $\nu^2 k^2 > 4c_s^2$  on the same plot (but a different plot from part d).