

Lecture 6 MHD Wave Observations and MHD Equilibria HwasD

I. Observations of MHD Waves:

A. Alfvén Waves in the Solar Wind (see Belcher, Davis & Smith, JGR, 74, 2302 (1969).)

1. Consider the eigenmodes for the Alfvén wave with $U_x = U_z = 0$, $U_y = U_0$

a. $(\omega^2 - k_{\parallel}^2 v_A^2) U_y = 0 \Rightarrow \omega = \pm k_{\parallel} v_A$

b. Note: For $\underline{k} = k_{\perp} \hat{x} + k_{\parallel} \hat{z}$, $\underline{k} \cdot \underline{U}_1 = 0$ (incompressible)

Thus $\rho_1 = 0$ and $p_1 = 0$

c. $\omega \underline{B}_1 = \underline{B}_0 (\underline{k} \cdot \underline{U}_1) - (\underline{B}_0 \cdot \underline{k}) \underline{U}_1 \Rightarrow \omega B_y = -B_0 k_{\parallel} U_y$


Thus $\frac{B_y}{B_0} = -\frac{k_{\parallel}}{\omega} U_y = -\frac{k_{\parallel}}{(\pm k_{\parallel} v_A)} U_0 = \mp \frac{U_0}{v_A}$

d. $\boxed{\frac{B_y}{B_0} = \mp \frac{U_0}{v_A}} \quad \boxed{U_y = U_0} \quad \boxed{\omega = \pm k_{\parallel} v_A}$


2. For the Alfvén wave, the fluctuation $\left(\frac{B_y}{B_0}\right) / \left(\frac{U_y}{v_A}\right) = \mp 1$

a. B_y & U_y are either in phase or π out of phase

3. Measurements of magnetic field fluctuations in solar wind



I (Continued)

B. Radio Emissions from Jupiter:

Proposed by:

Goldreich & Lynden-Bell

Astrophys. J., 156, 59, (1969).

1. The motion of Jupiter's moon Io through the plasma in Jupiter's magnetic causes Alfvén waves.
2. These Alfvén waves propagate along the magnetic field to the upper atmosphere of Jupiter
3. When waves strike the atmosphere, the decelerations due to the current associated with the Alfvén wave (remember $\mathbf{J} = \frac{1}{\mu_0} \nabla \times \mathbf{B}$) cause radio emission.
4. Confirmed during Voyager I flyby of Io [see Ness, ^{NFI} et al. Science, 204, 982 (1979).]

II. MHD Equilibrium

A. Basic concepts about MHD Equilibrium

1. A static equilibrium in MHD exists when $\underline{U} = 0$ and the forces on the plasma are in balance.


2. Momentum (or force) Equation Force per unit volume = $\rho \frac{d\underline{U}}{dt} + \rho \underline{U} \cdot \nabla \underline{U} = -\nabla p + \underline{j} \times \underline{B} - \rho \nabla \Phi_G$

3. A static equilibrium satisfies $\boxed{0 = -\nabla p + \underline{j} \times \underline{B} - \rho \nabla \Phi_G}$

NOTE: Equilibrium here refers to force balance, not to thermal equilibrium.

B. Example: Hydrostatic Equilibrium in a plane atmosphere

1. In the earth's atmosphere, the gravitational force on a fluid can often be well approximated by $-\rho \nabla \Phi_G = -\rho g \hat{z}$



2. Thus the hydrostatic equilibrium gives

a. $0 = -\nabla p - \rho g \hat{z}$

b. $\Rightarrow \boxed{\frac{\partial p}{\partial z} = -\rho g}$ (z-component)

3. Consider an Isothermal Atmosphere with $T = T_0$.

a. The ideal gas law gives $p = nkT_0 = \frac{\rho k T_0}{m} \Rightarrow \rho = \frac{m}{k T_0} p$

4. Thus $\frac{\partial p}{\partial z} = -\frac{gm}{kT_0} p \Rightarrow$ Solution $p(z) = p_0 e^{-\frac{z}{H_0}}$

where the scale height ~~is~~ $H_0 = \frac{kT_0}{gm}$.

5. The pressure and density in an isothermal atmosphere vary exponentially with scale height $H_0 = \frac{kT_0}{gm}$.

II. (Continued)

C. Back to MHD

1. If gravity is negligible (often true in the hot, diffuse plasmas in laboratory devices), then the MHD force balance is

$$\nabla p = \underline{j} \times \underline{B}$$

a. NOTE: We naturally must also satisfy Maxwell's Eqs: $\nabla \times \underline{B} = \mu_0 \underline{j}$
 $\nabla \cdot \underline{B} = 0$

2. Taking the dot product of the force balance eq. with \underline{j} or \underline{B} gives

$$\underline{j} \cdot \nabla p = 0$$

$$\underline{B} \cdot \nabla p = 0$$


Thus, in force balance \underline{B} and \underline{j} must lie on surfaces of constant pressure.

3. Magnetic Fusion: The goal of fusion is to contain a hot plasma with a magnetic field.

a. In Ideal MHD, the Frozen-In Flux Theorem tells us that the magnetic field \underline{B} is frozen to the plasma.

b. Vice versa, the plasma is stuck to the magnetic field lines.

c. BUT, the plasma can move freely along the \underline{B} field line.



d. So, to confine a hot plasma, surfaces of constant pressure must be closed.

4. Static Equilibria can be divided into two classes:

| | |
|---|----------------|
| $\underline{j} \times \underline{B} = 0$ | Force-free |
| $\underline{j} \times \underline{B} \neq 0$ | Force-Balanced |

III. Force-Free MHD Equilibrium $\underline{j} \times \underline{B} = 0$

1. This can occur in tenuous or cold plasmas where the pressure can be neglected.

2. The condition $\underline{j} \times \underline{B} = 0$ is always satisfied when $\mu_0 \underline{j} = \nabla \times \underline{B} = \alpha \underline{B}$ where α is a scalar function.

3. What does the force free condition satisfied by $\nabla \times \underline{B} = \alpha \underline{B}$ tell us?

a. NOTE! Taking the ~~curl~~ divergence

$$\nabla \cdot (\nabla \times \underline{B}) = \nabla \cdot (\alpha \underline{B})$$

$$0 = \alpha (\nabla \cdot \underline{B}) + \underline{B} \cdot \nabla \alpha \Rightarrow \underline{B} \cdot \nabla \alpha = 0$$

b. Again, \underline{B} ~~must~~ field lines must lie on surfaces of constant α .

4. In this case, we can show

The constant α surface cannot be a simple, closed surface.

Proof: a. Let C be a closed curve along a magnetic field line \underline{B} .

Thus $\int_C \underline{B} \cdot d\underline{l} \neq 0$

$$b. \int_C \underline{B} \cdot d\underline{l} = \int_S (\nabla \times \underline{B}) \cdot d\underline{A} = \int_S \alpha \underline{B} \cdot d\underline{A}$$

Stoke's Thm
NRL PB (34)

$$\nabla \times \underline{B} = \alpha \underline{B}$$

Surface bounded by
Curve C.

c. Displace surface S until it coincides with a surface of const α .
This can only be done if the surface is simply closed
(otherwise Stoke's theorem is invalid)

d. Since $\alpha = \text{const}$, we have $= \alpha \int_S \underline{B} \cdot d\underline{A}$

II (Continued)
Axi. (Continued)

e. But, \underline{B} lines on the surface of constant α , so


$$\int_S \underline{B} \cdot d\underline{A} = 0.$$

Ⓐ We have reached a contradiction!

Ⓑ Thus, the surface of constant α cannot be a simply closed surface! QED!

Ⓒ


5. A spherical surface of constant α will not work!



a. The magnetic field on a simple closed surface cannot be divergence free

"You can't comb the hair everywhere."

6. Hopf's Theorem: The simplest topological surface of constant α that satisfies both conditions $\nabla \cdot \underline{B} = 0$ and $\underline{B} \cdot \nabla \alpha = 0$ is a torus.



a. This theorem also holds for force-balance equilibria ($\nabla p \neq 0$). Here the surfaces are constant pressure surfaces.

B. Flux Ropes:

cylindrical geometry

Cylindrical
(r, ϕ, z)

↳ $\nabla \times \underline{B} = \alpha \underline{B}$ can be satisfied by $\underline{B} = B_\phi \hat{\phi} + B_z \hat{z}$ with

$$B_\phi = \frac{B_0 k r}{1 + k^2 r^2}$$

$$B_z = \frac{B_0}{1 + k^2 r^2}$$

Lecture 6 (Continued)

II. B (Continued)

2. The z -component of the current $(j_z = \frac{1}{\mu_0} \nabla \times B_z)$ is then

$$\text{and } \alpha = \frac{\mu_0 j_z}{B_z} = \frac{2k}{1+k^2 r^2}$$


$$j_z = \frac{2k B_0}{\mu_0 (1+k^2 r^2)^2}$$

Strong current dies off $\propto \frac{1}{r^4}$

Howes ⑦

3. Geometry of Flux Ropes:


- B consists of nested helices.
- Pitch angles (w.r.t. z) from 0 at axis ($r=0$) to $\frac{\pi}{2}$ at large r .



pitch angle increases with $r \rightarrow \lim_{r \rightarrow \infty} \frac{B_\theta}{B_z} \propto r$

4. Observations:

- Prominences in the solar corona can be seen to have such a helical structure.



C. Reverse Field Pinch (RFP)

1. Another class of force-free solutions with cylindrical symmetry have

$$B_\theta = B_0 J_1(\alpha r)$$

$$B_z = B_0 J_0(\alpha r)$$


a. α, B_0 are positive constants

b. J_0 & J_1 are zeroth & first order Bessel functions.

III. C. (Continued)

2. ~~Plasma~~ Magnetic Fields again are nested helices with varying pitch angle.


3. For a plasma of radius $a > \frac{2.4}{\alpha}$, the axial field reverses outside the first zero of the Bessel function J_0 .



Such a configuration is known as a Reverse Field Pinch (RFP)

These have good confinement properties for fusion.

$\alpha r = 2.4 \Rightarrow J_0(\alpha r) = 0$



Surface with $B_z = B_0 J_0(\alpha r) = 0$

D. Taylor Relaxation [Taylor, J.B. Phys. Rev. Lett. 33, 1139 (1974)]

1. Taylor proposed RFP's spontaneously form when a turbulent toroidally confined plasma is allowed to relax

a. Magnetic Helicity $H_m \equiv \int d^3x \underline{A} \cdot \underline{B}$ remains constant (ie is a conserved quantity in IDEAL MHD) while

b. Plasma relaxes to state of minimum potential energy

Energy $E = \frac{1}{2} \rho v^2 + \frac{P}{\gamma - 1} + \frac{B^2}{2\mu_0}$

Kinetic Energy Potential Energy.

Taylor Relaxation