

Lecture H21: The Physics of Accretion

I. The Physics of Accretion

Ref: Frank, King, & Raine, Accretion Power in Astrophysics, Cambridge Univ Press, 2002.

A. History

1. 19th century: Gravity was known source of astrophysical energy, but was inadequate to power the sun \rightarrow nuclear energy!
2. Present: The most luminous objects in the universe (AGN, quasars, etc) cannot be powered by nuclear burning \rightarrow accretion power is the critical driver of these systems!

B. Nuclear vs. Accretion Power

i. Estimate of energy gain in accretion

a. Gravitational Potential $\Phi_G = -\frac{GM}{r}$

b. For an object of mass M and radius R_* , the gravitational potential energy released as an object falls from $r = \infty$ is

$$\Delta E_{\text{acc}} = -\left(\frac{-GMm}{R_*} - \frac{-GMm}{\infty}\right) = \frac{GMm}{R_*} = \boxed{\Delta E_{\text{acc}}}$$

2. Estimates for typical objects:

a. Sun: $M = M_\odot = 2 \times 10^{33} \text{ g}$ $R_\odot = 7 \times 10^{10} \text{ cm}$ $G = 6.67 \times 10^{-8} \frac{\text{dyne cm}^2}{\text{g}^2}$

i. $\frac{\Delta E_{\text{acc}}}{m} = \frac{GM_\odot}{R_\odot} = \frac{(6.67 \times 10^{-8} \frac{\text{dyne cm}^2}{\text{g}^2})(2 \times 10^{33} \text{ g})}{(7 \times 10^{10} \text{ cm})} = 2 \times 10^{15} \frac{\text{ergs}}{\text{g}}$

ii. NOTE: We will adopt cgs units for consistency with astrophysical convention.

iii. For comparison, $L_\odot = 3.83 \times 10^{33} \frac{\text{cm}}{\text{s}}$

- This requires $\dot{M} = \frac{L_\odot}{\Delta E_{\text{acc}}/m} = \frac{3.83 \times 10^{33} \frac{\text{cm}}{\text{s}}}{2 \times 10^{15} \frac{\text{ergs}}{\text{g}}} = 2 \times 10^{18} \frac{\text{g}}{\text{s}}$

- Accretion of the entire solar mass would take $T = \frac{M}{\dot{M}} = \frac{2 \times 10^{33} \text{ g}}{2 \times 10^{18} \frac{\text{g}}{\text{s}}} = 10^{15} \text{ s} \approx 10^{15} \text{ s} / (3.15 \times 10^7 \text{ s}) = 3.0 \times 10^7 \text{ yr} = 30 \text{ My}$

Lecture #21 (Continued)

Hanes (3)

Z.B. 2.a.iii (Continued)

- Therefore, since $T \sim 30 \text{ Myr} \ll$ lifetime of the sun, accretion is insufficient to power the sun.

b. White Dwarf: $M \sim M_\odot$, $R_* \sim 10^9 \text{ cm}$

$$\frac{\Delta E_{\text{acc}}}{m} = \frac{GM_\odot}{R_*} = \frac{(6.67 \times 10^{-8} \frac{\text{dyn cm}^2}{\text{g}^2})(2 \times 10^{33} \text{ g})}{10^9 \text{ cm}} = 10^{17} \frac{\text{erg}}{\text{g}}$$

c. Neutron Star: $M \sim M_\odot$, $R_* \sim 10 \text{ km} \sim 10^6 \text{ cm}$

$$\frac{\Delta E_{\text{acc}}}{m} = \frac{GM_\odot}{R_*} = \frac{(6.67 \times 10^{-8} \frac{\text{dyn cm}^2}{\text{g}^2})(2 \times 10^{33} \text{ g})}{10^6 \text{ cm}} \sim 10^{20} \frac{\text{erg}}{\text{g}}$$

d. Black Hole: $M \sim M_\odot$, $R_* \sim \frac{2GM}{c^2} \sim 3 \times 10^5 \text{ cm}$

$$\frac{\Delta E_{\text{acc}}}{m} \sim \frac{GM_\odot}{R_*} \sim \frac{GM_\odot}{\left(\frac{2GM}{c^2}\right)} \sim \frac{c^2}{2} \sim 5 \times 10^{20} \frac{\text{erg}}{\text{g}} \rightarrow \Delta E_{\text{acc}} \sim \frac{1}{2} mc^2$$

Half-remnant energy.

- Note, however, that energy released from gravitational potential may pass through event horizon and not be emitted (more later).

3. Nuclear Burning Energy Release

a. Fusion of hydrogen to helium releases an energy

$$\Delta E_{\text{nuc}} = 0.007 mc^2$$

b. Thus $\frac{\Delta E_{\text{nuc}}}{m} \sim 6 \times 10^{18} \frac{\text{erg}}{\text{g}}$

c. a. Nuclear burning is much less efficient than accretion for compact objects such as neutron stars or black holes.

b. $\Delta E_{\text{nuc}} \sim 60 \Delta E_{\text{acc}}$ for white dwarfs, but nuclear burning is episodic

c. $\Delta E_{\text{nuc}} \gg \Delta E_{\text{acc}}$ for the Sun.

d. Compaction of object is important for accretion to be dominant!

Leave #2 (Continued)

Hawes(3)

I.C. The Eddington Limit

1. Here, we consider the limiting luminosity due to accretion if radiation pressure balances gravitational force.

2. Consider spherically symmetric accretion, and Steady flow.

3. For a hydrogenic (proton plus electron) fully ionized plasma, radiation exerts a force mainly on free electrons through

Thompson Scattering, $\sigma_T = 6.7 \times 10^{-25} \text{ cm}^2$

a. For a radiative energy flux $S (\frac{\text{erg}}{\text{s cm}^2})$, momentum from scattering is

$$\frac{\sigma_T S}{c}$$

b. This pushes out on electrons, but plasma waves to maintain quasineutrality, so the electrons drag the protons with them.

4. Gravity: $\frac{GM(m_p + m_e)}{r^2} \approx \frac{GMm_p}{r^2}$

5. Balance Gravity and radiation pressure force.

a. $S = \frac{L}{4\pi r^2}$ for an accreting source of Luminosity $L (\frac{\text{erg}}{\text{s}})$

b. $\frac{GMm_p}{r^2} - \frac{L\sigma_T}{4\pi c r^2} = 0$

c. $L_{\text{edd}} = \frac{4\pi GMm_p c}{\sigma_T} = 1.3 \times 10^{38} \left(\frac{M}{M_\odot}\right) \frac{\text{erg}}{\text{s}} = L_{\text{edd}}$

d. This gives the maximum Luminosity for a steady, spherically symmetric accretion powered object.

e. Note that unsteady events (supernova explosions) can significantly exceed this limit. ($L \sim 10^{42} \text{ to } 10^{43} \frac{\text{erg}}{\text{s}}$ at peak)

Lecture 21 (Continued)

Answers (4)

7. C. (Continued)

6. Accretion Luminosity:

$$a. L_{\text{acc}} = \frac{GM\dot{M}}{R_*}$$

$$b. \text{White Dwarfs: } L_{\text{acc}} = 1.3 \times 10^{33} \dot{M}_{16} \left(\frac{M}{M_\odot} \right) \left(\frac{10^9 \text{ cm}}{R_*} \right) \frac{\text{erg}}{\text{s}}$$

$$\text{where } \dot{M}_{16} = \frac{\dot{M}}{10^{16} \text{ g/s}} \quad \text{Typical } \dot{M} \text{ in close binary systems.}$$

$$c. \text{Neutron Star: } L_{\text{acc}} = 1.3 \times 10^{36} \dot{M}_{16} \left(\frac{M}{M_\odot} \right) \left(\frac{10 \text{ km}}{R_*} \right) \frac{\text{erg}}{\text{s}}$$

7. Black Holes:

a. Since black hole can swallow some energy released in accretion, we model this by an efficiency, η .

$$L_{\text{acc}} = \frac{2\eta GM\dot{M}}{R_*} = \frac{2\eta GM\dot{M}}{(2GM/c)} = \eta \dot{M}c^2$$

↑ efficiency for converted accreted rest mass to radiation.

b. Typical values from modeling are $\eta \sim 0.1$

D. Spectrum of Accretion Powered Emission

1. Characterize frequency of emitted radiation: $h\nu \sim T_{\text{rad}}$

photon energy \sim thermal energy

2. Black body radiation:

$$a. S = \frac{1}{4\pi r^2} = \sigma T^4 \quad \sigma = 5.67 \times 10^{-5} \frac{\text{erg}}{\text{cm}^2 \text{s} \text{K}^4}$$

$$b. \text{Thus } T_b = \left(\frac{L_{\text{acc}}}{4\pi R_*^2 \sigma} \right)^{\frac{1}{4}} \quad \text{"Blackbody temperature"}$$

3. Thermal Temperature: a. Potential energy: $\frac{GM(\text{infinite})}{R_*} = \frac{GM_{\text{mp}}}{R_*}$

b. Thermal energy: $\frac{2}{3} \times \frac{3}{2} T \approx 3T$
ions & electrons

$$c. \text{Thus, } T_{\text{fh}} = \frac{GM_{\text{mp}}}{3R_*}$$

Lecture #21 (Continued)

Hawes(5)

I. D. (Continued)

4. Conversion of accretion energy to radiation: Two limits.

a. In optically thick conditions, radiation reaches thermal equilibrium with radiated material before escaping, so $T_{\text{rad}} \sim T_b$

b. If accretion energy is directly converted to radiation, $T_{\text{rad}} \sim T_{\text{th}}$.

c. In general, we have

$$T_b \leq T_{\text{rad}} \leq T_{\text{th}}$$

5. a. Neutron Star

$$L_{\text{acc}} \sim 10^{36} \frac{\text{erg}}{\text{s}}$$

$$1 \text{ keV} \leq h\nu \leq 50 \text{ MeV}$$

(X-rays)

(γ -rays)

b. White Dwarf

$$L_{\text{acc}} \sim 10^{33} \frac{\text{erg}}{\text{s}}$$

$$6 \text{ eV} \leq h\nu \leq 100 \text{ keV}$$

(UV)

(X-rays)

6. Electromagnetic Spectrum	Classification	Wavelength (nm)	Energy
	X-rays	≤ 0.01	$\gtrsim 100 \text{ keV}$
	Hard X-rays	$0.01 - 1$	$1 - 100 \text{ keV}$
	Soft X-rays	$1 - 10$	$100 \text{ eV} - 1 \text{ keV}$
	Extreme UV	$10 - 121$	$10 - 100 \text{ eV}$
	UV	$122 - 400$	$3 - 10 \text{ eV}$
	Visible	$400 - 760$	$\leq 3 \text{ eV}$

II. Bondi Accretion:

Ref: Bondi (1952) 'On spherically symmetrical accretion', MNRAS 112, 195.

A. Motivation.

- For spherically symmetric accretion onto a star at rest with respect to the interstellar medium (ISM), can we determine the accretion rate and effective radius in terms of stellar mass and ISM properties?
- We'll see this is essentially the same as the Parker-Sohr Wind Solution.

Lecture # 21 (Continued)

Howes ⑥

II. B. Bondi Accretion Solution

1. Assumptions: a. Spherically Symmetric, $\frac{\partial}{\partial \phi} = 0$, $\frac{\partial}{\partial \theta} = 0$

b. Steady-state, $\frac{\partial}{\partial t} = 0$

c. Radial inflow, $\underline{U} = U_r(r) \hat{r}$, $U_\phi = U_\theta = 0$

d. Ignore Magnetic Field, $\underline{B} = 0$

2 MHD Equations: a. Continuity: $\frac{\partial p}{\partial r} + \nabla \cdot (\rho \underline{U}) = 0$

b. Momentum: $\rho \frac{\partial \underline{U}}{\partial r} + \rho (\underline{U} \cdot \nabla) \underline{U} = -\nabla p + \frac{\rho \underline{B} \times \underline{B}}{4\pi} - \rho \frac{GM}{r^2} \hat{r}$

c. Induction: $\underline{B} = 0$, so no required

d. Polytropic Relation: $p = k \rho^\gamma$ [same as $\frac{d}{dr} \left(\frac{p}{\rho^\gamma} \right) = 0$]

3. NOTE: Sound speed $C_s^2 = \left(\frac{dp}{d\rho} \right)$

a. Adiabatic ($\gamma = \frac{5}{3}$):

$$C_s = \left(\frac{\gamma p}{\rho} \right)^{\frac{1}{2}} \propto \rho^{\frac{1}{3}}$$

b. Isothermal ($\gamma = 1$):

$$C_s = \left(\frac{k}{\rho} \right)^{\frac{1}{2}} = \text{constant.}$$

4. Ideal Gas Law: $p = \frac{\rho T}{\mu m_p}$, where μ is mean molecular weight.

a. For hydrogen $\mu = 1$

b. For ionized hydrogen (protons+electrons), $\mu = \left(\frac{m_p + m_e}{2} \right) \frac{1}{2} \approx \frac{m_p}{2m_p} = \frac{1}{2}$

c. Thus, for a fully ionized plasma, $\boxed{p = \frac{2\rho T}{m_p}}$ (same as before, Lect 18)

5. Continuity Equation: $\frac{1}{r^2} \frac{\partial}{\partial r} (r^2 \rho U_r) = 0 \Rightarrow \boxed{4\pi r^2 \rho U_r = \dot{M}}$

Mass accretion rate

6. Radial Momentum: $\rho U_r \frac{\partial U_r}{\partial r} + \frac{\partial p}{\partial r} + \rho \frac{GM}{r^2} = 0$

a. $\frac{\partial p}{\partial r} = \left(\frac{\partial p}{\partial \rho} \right) \frac{\partial \rho}{\partial r} = C_s^2 \frac{\partial \rho}{\partial r}$

b. Thus, $U_r \frac{\partial U_r}{\partial r} + \frac{1}{\rho} \left[C_s^2 \frac{\partial \rho}{\partial r} \right] + \frac{GM}{r^2} = 0$

c. From continuity eq., $r^2 U_r \frac{\partial U_r}{\partial r} + \rho \frac{\partial}{\partial r} (r^2 U_r) = 0 \Rightarrow \frac{1}{\rho} \frac{\partial \rho}{\partial r} = -\frac{1}{r^2 U_r} \frac{\partial}{\partial r} (r^2 U_r)$

Lecture #21 (Continued)

Hawes 7

II.B.6. (Continued)

d. This $U_r \frac{\partial U_r}{\partial r} - \frac{C_s^2}{r^2 U_r} \frac{\partial(r^2 U_r)}{\partial r} + \frac{GM}{r^2} = 0$

$$= r^2 \frac{\partial U_r}{\partial r} + 2r U_r$$

e. Finally

$$\frac{1}{2} \left(1 - \frac{C_s^2}{U_r^2} \right) \frac{\partial(U_r^2)}{\partial r} = - \frac{GM}{r^2} \left[1 - \frac{2C_s^2 r}{GM} \right]$$


7. This is the same as the Parker-Sobr wind solution with general γ :

a. Critical radius: $r_s = \frac{GM}{2C_s^2} \approx 7.5 \times 10^{13} \left(\frac{I}{10^4 K} \right)^{-1} \left(\frac{M}{M_\odot} \right) \text{ cm}$

using $C_s \approx 10 \left(\frac{I}{10^4 K} \right)^{1/2} \frac{\text{km}}{\text{s}}$ as sound speed for $1 \leq \gamma \leq \frac{5}{3}$

b. At $r = r_s$, LHS = 0 requires i) $U_r^2 = C_s^2$
or ii) $\frac{\partial U_r^2}{\partial r} = 0$

8. Solutions:



Four Types of Solutions

- a) Unphysical Solutions ⑤ & ⑥
- b) Purely Supersonic ④
- c) Purely Subsonic ③
- d) Transonic Solutions ① & ②

9. a) We expect $U_r \rightarrow 0$ as $r \rightarrow \infty$ for acceleration problem \rightarrow Solutions ① & ③
- b) Since RHS < 0 for small $r \rightarrow 0$, this implies $U_r > C_s$ as $r \rightarrow 0 \Rightarrow$ Solution ①!

Lecture #21 (Continued)

Howes ⑧

IB. 9. c. Solution ① is unique solution for accretion problem!

C. Properties of Bondi Accretion Flow:

1. Integrate ① on $U_r \frac{\partial U_r}{\partial r} + \frac{1}{\rho} \frac{\partial p}{\partial r} + \frac{GM}{r^2} = 0$

a. $\frac{U_r^2}{2} + \int \frac{dp}{\rho} - \frac{GM}{r} = \text{const.}$

b. NOTE: $dp = \gamma K p^{\gamma-1} dp$, so $\int \frac{dp}{\rho} = \gamma K \int \frac{p^{\gamma-1}}{\rho} dp = \frac{\gamma K}{\gamma-1} p^{\gamma-1}$
 $= \frac{\gamma K p^\gamma}{(\gamma-1) \rho} = \frac{1}{\gamma-1} \left(\frac{\gamma p}{\rho} \right) = \frac{C_s^2}{\gamma-1}$

c. Thus, we obtain the Bernoulli Integral

$$\boxed{\frac{U_r^2}{2} + \frac{C_s^2}{\gamma-1} - \frac{GM}{r} = \text{const}}$$

d. At $r \rightarrow \infty$, $U_r \rightarrow 0$, so $\frac{U_r^2}{2} + \frac{C_{s\infty}^2}{\gamma-1} - \frac{GM}{r} = \text{const} \leftarrow \text{constant determined.}$

$$\boxed{\frac{U_r^2}{2} + \frac{C_s^2}{\gamma-1} - \frac{GM}{r} = \frac{C_{s\infty}^2}{\gamma-1}}$$

2. At $r = r_s$, solution ① gives $U_r^2 = C_s^2(r_s)$, so $C_s^2(r_s) \left[\frac{1}{2} + \frac{1}{\gamma-1} - 2 \right] = \frac{C_{s\infty}^2}{\gamma-1}$

c. Thus $\boxed{C_s(r_s) = C_{s\infty} \left(\frac{2}{5-3\gamma} \right)^{\frac{1}{2}}}$ connects sound speed at orbital radius to sound speed in ZSM, $C_{s\infty}$
 (Valid for $1/\gamma < \frac{5}{3}$)

3. Connect $\rho(r_s)$ to ρ_∞ :

a. $C_s^2 \times \frac{\partial p}{\rho} \propto p^{\gamma-1} \Rightarrow \frac{C_s^2(r_s)}{C_{s\infty}^2} = \left(\frac{\rho(r_s)}{\rho_\infty} \right)^{\gamma-1} \rightarrow \boxed{\rho(r_s) = \rho_\infty \left[\frac{C_s(r_s)}{C_{s\infty}} \right]^{\frac{2}{\gamma-1}}}$

4. Obtain mass accretion rate \dot{M} in terms of $C_{s\infty}$ & ρ_∞ :

a. $\dot{M} = 4\pi r_s^2 \rho U_r = 4\pi r_s^2 \rho(r_s) C_s(r_s)$

b Substituting for r_s , $\rho(r_s)$, & $C_s(r_s)$ in terms of $C_{s\infty}$ & ρ_∞ ,

Lesson #2.1 (Continued)

Hawes ⑨

II.C. 4. (Continued)

c. $\dot{M} = \pi G^2 M^2 \frac{\rho_{\infty}}{C_{S\infty}^3} \left(\frac{2}{5-3\gamma} \right)^{\frac{5-3\gamma}{2(\gamma-1)}}$ for $1 < \gamma < \frac{5}{3}$

d. Typical Values $C_{S\infty} = 10 \frac{\text{km}}{\text{s}}$ and $\rho_{\infty} = 10^{-24} \frac{\text{g}}{\text{cm}^3}$ ($n \approx 1 \text{ cm}^{-3}$)

$$\dot{M} \approx 1.4 \times 10^{11} \left(\frac{M}{M_{\odot}} \right)^2 \left(\frac{\rho_{\infty}}{10^{-24} \frac{\text{g}}{\text{cm}^3}} \right) \left(\frac{C_{S\infty}}{10 \frac{\text{km}}{\text{s}}} \right)^{-3} \frac{\text{g}}{\text{s}}$$
 for $\gamma = 1.4$

This accretion rate is quite low, and Bondi accretion powered radiation is probably not observable ($L_{\text{acc}} \sim 10^{31} \frac{\text{erg}}{\text{s}}$ for neutron star).

NOTE: For binary mass transfer, typically $\dot{M} \sim 10^6 \frac{\text{g}}{\text{s}}$ (much larger)

5. Accretion radius, r_{acc} :

a. Bernoulli Integral: $\frac{U_r^2}{2} + \frac{C_s^2 - C_{S\infty}^2}{\gamma - 1} - \frac{GM}{r} = 0$

b. At $r \rightarrow \infty$, gravity is weak, $U_r \rightarrow 0$, and $C_s = C_{S\infty}$.

c. As r decreases, inflow velocity reaches $U_r = -C_{S\infty}$ (inflow)
- Only gravitational term can balance this flow term, so

$$\frac{U_r^2}{2} \approx \frac{GM}{r_{\text{acc}}} \Rightarrow r_{\text{acc}} \approx \frac{2GM}{C_{S\infty}^2}$$

d.

$$r_{\text{acc}} = 3 \times 10^{14} \left(\frac{M}{M_{\odot}} \right) \left(\frac{10^4 K}{T_{\infty}} \right) \text{ cm}$$

0. Summary of Bondi Accretion Flow:

1. Accretion Rate: $\dot{M} = \pi G^2 M^2 \frac{\rho_{\infty}}{C_{S\infty}^3} \left(\frac{2}{5-3\gamma} \right)^{\frac{5-3\gamma}{2(\gamma-1)}}$

2. Accretion Radii: $r_{\text{acc}} = \frac{2GM}{C_{S\infty}^2}$

3. Accretion from ISM to isolated stars is unlikely to be observable
 \Rightarrow Must look at binaries to find more powerful accreting systems.