

Hawes D

Lecture #21 The Structure of the Heliosphere

I. Solar and Stellar Winds:

Refs.: 1. Weber & Davis, 1967, ApJ. 148, 127.

2. Lomas & Cassinelli, Introduction to Stellar Winds, Cambridge Univ. Press, 1999.

A. Equatorial Stellar Wind Model

1. Consider a model for the Solar wind in spherical coords, (r, θ, ϕ)

$$\mathbf{U} = U_r(r) \hat{r} + U_\theta(r) \hat{\theta}$$

$$\mathbf{B} = B_r(r) \hat{r} + B_\theta(r) \hat{\theta}$$

2. By symmetry on equatorial plane, $U_\theta = 0$ and $B_\theta = 0$.

3. Assume Steady State conditions, $\frac{\partial}{\partial t} = 0$

4. From MHD, let's determine the global structure of the Interplanetary Magnetic Field (IMF) emerging from the rotating Sun with the solar wind flow.

5. Conservation of Mass:

a. Continuity Equation: $\frac{\partial \rho}{\partial t} + \nabla \cdot (\rho \mathbf{U}) = 0$

b. In spherical coordinates, $\frac{1}{r^2} \frac{\partial}{\partial r} (r^2 \rho U_r) = 0$

c. Thus $r^2 \rho U_r = \text{constant}$ ①

$$4\pi r^2 \rho U_r = \dot{M}$$

where $\dot{M} = \frac{\text{Mass Flux}}{\text{time}}$

6. Zero Magnetic Divergence:

a. $\nabla \cdot \mathbf{B} = 0$

b. Thus, $\frac{1}{r^2} \frac{\partial}{\partial r} (r^2 B_r) = 0$

c. So, $r^2 B_r = \text{constant}$

$$B_r = B_0 \left(\frac{R_0}{r} \right)^2 \quad \text{②}$$

B_0 is radial magnetic field at surface = R_0

Lecture #21 (Continued)
IA (Continued)

Haves (2)

7. Magnetic Induction Equations:

a. $\frac{\partial \mathbf{B}}{\partial t} = \nabla \times (\mathbf{U} \times \mathbf{B})$

b. Consider the θ -component of curl: $\frac{1}{r} \frac{\partial}{\partial r} [r(\mathbf{U} \times \mathbf{B})_\theta] = 0$

c. Thus, $\frac{1}{r} \frac{\partial}{\partial r} [r(U_r B_\phi - U_\phi B_r)] = 0$

d. $r(U_r B_\phi - U_\phi B_r) = \text{constant}$

At solar surface, $U_\phi \gg U_r$, and $U_\phi = R_\odot \Omega_\odot$, so

constant = $-R_\odot^2 \Omega_\odot B_\phi$ where $\Omega_\odot = 2.85 \times 10^{-6} \frac{\text{rad}}{\text{s}}$

e. Since $r^2 B_r = R_\odot^2 B_\phi$, we obtain constant = $-r^2 \Omega_\odot B_r$

Substituting this result above and simplifying,

$$\boxed{\frac{B_\phi}{B_r} = \frac{U_\phi - r \Omega_\odot}{U_r}} \quad (3)$$

8. Conservation of Angular Momentum:

a. $\rho \frac{\partial \mathbf{U}}{\partial t} + \rho \mathbf{U} \cdot \nabla \mathbf{U} = -\nabla(p + \frac{B^2}{2\mu_0}) + \frac{\mathbf{B} \cdot \nabla \mathbf{B}}{\mu_0} - \rho \frac{GM}{r^2} \hat{r}$

b. The azimuthal component gives, in steady state & spherical coords,

$$\rho U_r \frac{\partial}{\partial r} (r U_\phi) = \frac{B_r}{\mu_0} \frac{\partial}{\partial r} (r B_\phi)$$

c. $\frac{\partial}{\partial r} (r U_\phi) - \underbrace{\frac{B_r}{\mu_0 U_r} \frac{\partial}{\partial r} (r B_\phi)}_{\text{from DE (3)}} = 0$

From DE (3), $\frac{B_r}{\mu_0 U_r} = \left(\frac{4\pi}{M}\right) \left(\frac{B_\phi R_\odot^2}{\pi}\right) = \frac{4\pi B_\phi R_\odot^2}{M} = \text{constant}$

d. $\frac{\partial}{\partial r} \left(r U_\phi - \frac{r B_r B_\phi}{\mu_0 U_r} \right) = 0$

Lecture #21 (Continued)

Hawes (3)

I.A.8. (Continued)

e. $\boxed{L = r U_\phi - \frac{r B_r B_\phi}{\mu_0 \rho U_r}}$

(4)

Conservation of Angular Momentum,
where $L = \frac{\text{Angular Momentum}}{\text{unit mass}}$

9. Therefore we have

a. $4\pi r^2 \rho U_r = \dot{M}$

$$B_r = B_0 \left(\frac{R_0}{r} \right)^2$$

$$\frac{B_0}{B_r} = \frac{U_r - r \Omega_0}{U_r}$$

$$L = r U_\phi - \frac{r B_r B_\phi}{\mu_0 \rho U_r}$$

b. If $U_r(r)$ is known, we can solve for $\rho(r)$, $B_r(r)$, $B_\phi(r)$, and $U_\phi(r)$

c. Constants: $\dot{M}, B_0, \Omega_0, L$

c. We can continue, using Conservation of Energy and Radial Momentum Equation to obtain a fully self-consistent (and complicated) system, but for now we'll focus on the structure of B .

B. The Parker Spiral Magnetic Field

1. At distances $r \gg R_0$, $U_\phi \ll U_r$, so (3) can be simplified to

$$\frac{B_\phi}{B_r} \approx -\frac{r \Omega_0}{U_r}$$

2. Substituting (2), $B_\phi = -\frac{r \Omega_0 B_0 R_0^2}{U_r r^2} = -B_0 \frac{R_0^2 \Omega_0}{r U_r}$

3. Therefore, we see that $B_r \propto \frac{1}{r^2}$ and $B_\phi \propto \frac{1}{r}$, so the magnetic begins primarily radial and eventually becomes more and more azimuthal.

Lecture 21 (Continued)

Hoves (4)


I. B. (Continued)

4. One may obtain a function $\phi(r)$ that describes a magnetic field line,

$$\boxed{\phi(r) = \phi_0 - \frac{2\Omega}{V_{SW}}(r - R_A)} \quad \text{Archimedean Spiral}$$

where ϕ_0 is angle of "fase" of field line at $r = R_A$


and R_A is the critical radius for Alfvén waves, $R_A \approx 10 R_\odot$.



"Parker Spiral" of the
Interplanetary Magnetic Field.

4. Magnetic Field Magnitude, $B = (B_r^2 + B_\theta^2)^{1/2}$

$$\boxed{B = B_\odot \left(\frac{R_\odot}{r}\right)^2 \sqrt{1 + \frac{r^2 \Omega_\odot^2}{V_p^2}}}$$



C. Loss of Angular Momentum

1. Weber & Davis (1967) first demonstrated that the solar wind magnetic field

significantly enhanced the loss of angular momentum (spin down) of the Sun (in addition spin down by mass loss alone).


$$r \propto \frac{V_p}{\Omega_\odot} = 1 \text{ AU}$$

2. "Effective corotation" of solar wind with the Sun out to the Alfvén radius $R_A \approx 10 R_\odot$.

II. The Heliospheric Current Sheet

A. 1. Priestman & Kopp (1971) ²

computed a self-consistent
Steady state solution
of MHD equations with
Parker transonic solution
along each flow line.




2. Open field lines
from the poles

Eventually spread out to cover 4π solid angle beyond $\sim 3R_\odot$.

3. Along the equator, oppositely directed field lines from opposite poles come close together, leading to the Heliospheric Current Sheet required to support the jump in IMF field direction from Sunward to anti-sunward.

4. Because the magnetic and rotation axes of the Sun do not coincide, this current sheet becomes rippled, like a ballerina skirt.




a. When Earth's orbit passes through current sheet, it moves from a Magnetic Sector of sunward IMF to a sector of anti-sunward IMF.

II. B. Coronal Holes and Coronal Helmet Streamer Belt

1. The simplified picture above is reasonably accurate for solar minimum,

2. Coronal Holes:


low density regions
of open magnetic field
lines that appear
dark on coronagraphs.



3. Coronal Streamer Belt:

Bright teardrop shaped region (coronal helmet streamers) that appear to straddle field line reversals. Generally presumed to create a belt around the Sun where the heliospheric current sheet meets the corona, but may be more geoelectrically complicated during Solar maximum.

4. Connection to Solar Wind Speed:




a. Ulysses spacecraft went to high heliographic latitude, and at solar minimum found fast solar wind ($2700-8100 \text{ km/s}$) at high latitudes, and slow solar wind ($< 500 \text{ km/s}$) at low latitudes $\lesssim 30^\circ \text{ N/S}$.

b. Fast solar wind is assumed to originate at coronal holes

c. Slow solar wind is believed to originate at the after regions of coronal streamers.

II. Corotating Interaction Regions

A. i. Because the Sun is rotating, regions of fast solar wind can emerge from the Sun at a position radially behind previously emitted slow solar wind streams.



ii. The fast radial flow in fast streams catches up with and compresses the slow solar wind radially beyond it.

iii. Since the regions of fast & slow solar wind persist for several solar rotations, this pattern of compression and rarefaction repeats at each solar rotation.

iv. The region of compression are known as corotating interaction regions, or CIRs.

III. Global Structure of the Heliosphere

A. The Termination Shock

1. The supersonic solar wind at large heliocentric radius connects to a boundary condition $p \rightarrow 0$ at $r \rightarrow \infty$ (Lec #18)


2. But, if a finite pressure exists in the surrounding Local Interstellar Medium (ISM), the coronal expansion must eventually stop when solar wind pressure becomes smaller than ISM pressure.

Lecture A21 (Continued)

Hawes ⑧

II. A. (Continued)

3. A termination shock must develop to transition from supersonic flow to subsonic flow to enable the solar wind flow to stop.
4. Beyond this is the heliosopause, the tangential discontinuity separating plasma and magnetic field of Solar origin from that of the local ZSM.



5. Neptune's orbit is at $R \sim 30$ AU

6. a. Voyager I crossed the termination shock in the Northern heliospheric hemisphere at $R \sim 94$ AU in December 2004

b. Voyager II crossed the termination shock in the Southern hemisphere at $R \sim 84$ AU in August 2007.

7. The Interstellar Boundary Explorer (IBEX) mission, launched in October 2008, uses measurements of energetic neutral atoms to map the heliospheric boundaries, showing that the direction of the local ISM magnetic field B_{ISM} may influence the shape of the heliosphere.

8. From plasma oscillation measurements, Voyager I crossed heliospause on 25 Aug 2012

R/21/AU