


## Lecture #3: MHD Stability Analysis of Rayleigh-Taylor Instability

### I. Normal Mode Analysis

#### A. Setup



1. Density has exponential form in direction of gravity ( $\hat{z}$ )

$$\text{a. } \rho_0(x) = \rho_{00} e^{-\frac{x}{H}} \quad H = \text{Scale height of density}$$

b. For  $H > 0$ , density decreases with height (stable)

$H < 0$ , density increases with height (unstable)

2. For simplicity, we assume  $\vec{B}_z = 0$  (No variation along mean field).

NOTE: Such variations would bend the magnetic field lines, leading to magnetic tension (which stabilizes instability)

b. We also assume incompressible motion,  $\nabla \cdot \vec{U}_t = 0$

3. We want to analyze this problem for stability.

a. Normal Mode Analysis

b. Energy Principle

#### B. Using Linear Force Operator

1. We could use

$$-\rho_0 \omega^2 \xi = F(\xi)$$

to solve for the characteristic frequencies. But instead, we'll begin from equation of motion.

## Lecture #3 (Continued)

### 7.5 Using Equation of Motion:

#### 1. Momentum Eq:

$$\rho \frac{\partial \vec{U}}{\partial t} + \rho \vec{U} \cdot \nabla \vec{U} = -\nabla(p + \frac{B^2}{2\mu_0}) + \frac{(\vec{B} \cdot \nabla) \vec{B}}{\mu_0} + \rho \vec{g}$$

gravity, where  
 $\vec{g} = -g \hat{z}$

#### 2. Linearize above Equilibrium:

$$p = p_0(x) + \epsilon p_1(x)$$

$$\vec{U} = \vec{U}_0(x) + \epsilon \vec{U}_1(x)$$

$$\vec{B} = \vec{B}_0(x) \hat{z} + \epsilon \vec{B}_1(x)$$

$$p = p_0(x) + \epsilon p_1(x)$$

vertical  
direction  
(scalar)

vector

$$3. \text{Lowest Order: } O(1): \quad 0 = -\nabla(p_0 + \frac{B_0^2}{2\mu_0}) + p_0 \vec{g}$$

a. Equilibrium satisfies:

$$\frac{\partial}{\partial x} \left( p_0 + \frac{B_0^2}{2\mu_0} \right) = -p_0 g$$

Static  
Equilibrium

b. Notation:

$$p'_0 = \frac{\partial p_0}{\partial x}, \quad B'_0 = \frac{\partial B_0}{\partial x}, \quad p''_0 = \frac{\partial p_0}{\partial x}$$

$$\Rightarrow p''_0 + \frac{B'_0 B'_0}{\mu_0} = -p_0 g$$

#### 4. Next Order: $O(\epsilon)$ :

$$\rho \frac{\partial \vec{U}_1}{\partial t} = -\nabla(p_1 + \frac{B_0 \cdot B_1}{\mu_0}) + \frac{(\vec{B}_0 \cdot \nabla) \vec{B}_1}{\mu_0} + \frac{\vec{B}_1 \cdot \nabla \vec{B}_0}{\mu_0} + \rho_1 \vec{g}$$

①

②

③

④

⑤

a. Term ③  $(\vec{B}_0 \cdot \nabla) \vec{B}_1 = \vec{B}_0 \cdot \frac{\partial \vec{B}_1}{\partial x} = 0$

b. Term ④  $(\vec{B}_1 \cdot \nabla) \vec{B}_0 = \vec{B}_1 \cdot \frac{\partial \vec{B}_0}{\partial x} = \vec{B}'_1 \vec{B}_0 \hat{z}$

c. We can eliminate Term ② by taking the curl of this equation:

$$(\nabla \times \nabla \phi = 0)$$

scalar function.

Lecture #3 (Continued)  
 I.C. 4. (Continued)

Haves ③

d.  $\nabla \times \left( \rho_0 \frac{\partial \mathbf{U}_1}{\partial t} \right) = \nabla \times (B_0' B_x \hat{z}) + \nabla \times (-\rho_1 g \hat{x})$

e. Note: Since  $\frac{\partial}{\partial z} = 0$ ,  $\nabla = \frac{\partial}{\partial x} \hat{x} + \frac{\partial}{\partial y} \hat{y}$

f. Let's find the  $\hat{z}$ -component:

$$\begin{aligned} 1. \hat{z} \cdot \left[ \nabla \times \left( \rho_0 \frac{\partial \mathbf{U}_1}{\partial t} \right) \right] &= \frac{\partial}{\partial x} \left[ \rho_0 \frac{\partial U_y}{\partial t} \right] - \frac{\partial}{\partial y} \left[ \rho_0 \frac{\partial U_x}{\partial t} \right] \\ &= \rho_0' \frac{\partial U_y}{\partial t} + \rho_0 \frac{\partial^2 U_y}{\partial x \partial t} - \rho_0 \frac{\partial^2 U_x}{\partial y \partial t} \end{aligned}$$

2.  $\hat{z} \cdot \left[ \nabla \times (B_0' B_x \hat{z}) \right] = 0$

3.  $\hat{z} \cdot \left[ \nabla \times (-\rho_1 g \hat{x}) \right] = -\frac{\partial}{\partial x} [-\rho_1 g] = g \frac{\partial \rho_1}{\partial x}$

g. Thus, we find  $\boxed{\rho_0' \frac{\partial U_y}{\partial t} + \rho_0 \frac{\partial^2 U_y}{\partial x \partial t} - \rho_0 \frac{\partial^2 U_x}{\partial y \partial t} = g \frac{\partial \rho_1}{\partial x}}$

5. Fourier Transform in  $y$  and  $t$ :  $\tilde{U}_1(x) = \tilde{U}_1(x) e^{i(k_y y - \omega t)}$

a. Thus  $\frac{\partial}{\partial t} = -i\omega$        $\frac{\partial}{\partial y} = ik_y$

b. This yields

$$\boxed{\rho_0' \tilde{U}_Y + \rho_0 \frac{\partial \tilde{U}_Y}{\partial x} - \rho_0 i k_y \tilde{U}_X = -\frac{k_y g \rho_1}{\omega}}$$

6. We assume incompressible motion  $\nabla \cdot \mathbf{U}_1 = 0$

$$\boxed{\frac{\partial U_x}{\partial x} + i k_y U_Y = 0} \Rightarrow U_Y = \frac{i}{k_y} \frac{\partial U_x}{\partial x}$$

7. Continuity Equation  $\frac{\partial \rho}{\partial t} + \mathbf{U}_1 \cdot \nabla \rho + \rho_1 \nabla \cdot \mathbf{U}_1 = 0$

a.  $\mathcal{O}(\epsilon)$ :  $\frac{\partial A}{\partial t} + \tilde{U}_1 \cdot \nabla \rho_0 + \rho_0 \nabla \cdot \tilde{U}_1 = 0$

b.  $\boxed{-i\omega \rho_1 + U_X \rho_0' = 0} \Rightarrow \rho_1 = -\frac{i}{\omega} U_X \rho_0'$

Lecture #3 (Continued)  
L.C. (Continued)

Hawes (4)

8. Eliminate  $U_y$  &  $p_1$  in favor of  $U_x$ :

a. This yields:

$$\frac{\partial^2 U_x}{\partial x^2} + \frac{p_0'}{p_0} \frac{\partial U_x}{\partial x} - k_y^2 \left[ 1 + \frac{g}{\omega^2} \left( \frac{p_0'}{p_0} \right) \right] U_x = 0$$

9. Note: Since  $p_0(x) = p_{00} e^{-\frac{x}{H}}$ , we have  $\frac{p_0'}{p_0} = -\frac{1}{H}$

$$\frac{\partial^2 U_x}{\partial x^2} - \frac{1}{H} \frac{\partial U_x}{\partial x} - k_y^2 \left( 1 - \frac{g}{\omega^2 H} \right) U_x = 0$$

10. We can solve this with the help of an integrating factor.

a. Let  $U_x(x) = f(x) e^{\frac{x}{2H}}$

b. This gives  $\frac{\partial^2 U_x}{\partial x^2} - \frac{1}{H} \frac{\partial U_x}{\partial x} = \left( \frac{\partial^2 f}{\partial x^2} - \frac{f}{4H^2} \right) e^{\frac{x}{2H}}$

c. Thus, we find:

$$\frac{\partial^2 f}{\partial x^2} + \alpha^2 f = 0 \quad \text{where } \alpha^2 = k_y^2 \left( \frac{g}{H\omega^2} - 1 \right) - \frac{1}{4H^2}$$

11. The function  $f(x)$  must satisfy the boundary conditions

$$U_x(0) = U_x(a) = 0 \Rightarrow f(0) = f(a) = 0$$

a. General Solution:  $f(x) = f_0 \sin \alpha x + f_1 \cos \alpha x$

b.  $f(0) = f_0(0) + f_1(1) = 0 \Rightarrow f_1 = 0$ .

c.  $f(a) = f_0 \sin(\alpha a) = 0 \Rightarrow \alpha = \frac{n\pi}{a} \quad \text{for } n = 1, 2, 3, \dots$

d. We therefore have eigenfunctions  $f_n$  with mode number  $n$ .

12. Solve for frequency:  $\frac{n^2 \pi^2}{a^2} = k_y^2 \left( \frac{g}{H\omega_n^2} - 1 \right) = \frac{1}{4H^2}$

a.  $\frac{n^2 \pi^2}{a^2} + k_y^2 + \frac{1}{4H^2} = \frac{g}{H} \frac{k_y^2}{\omega_n^2}$

Lesson #3 (Continued)

I.C.2 (Continued)

Hawes 5

b.  $\omega_n^2 = \frac{g}{H} - \frac{4H^2 k_y^2 a^2}{a^2 + 4H^2 (h^2 \pi^2 + k_y^2 a^2)}$

$\underbrace{\phantom{4H^2 k_y^2 a^2}}_{\text{positive definite}}$

c. For  $H > 0$ ,  $\omega_n^2 > 0 \Rightarrow$  Oscillating function  $\Rightarrow$  STABLE  
For  $H < 0$ ,  $\omega_n^2 < 0 \Rightarrow$  UNSTABLE.

d. 1. Max growth rate  $\omega_n^2 = \frac{g}{H}$  as  $k_y \rightarrow \infty$

2. Growth rate  $\rightarrow 0$  as  $k_y \rightarrow 0$

3. Lower vertical mode numbers  $n$  have faster growth.

## II. Energy Principle:

### A. Gravitational Force Term:

1. In Linear Force Operator  $\tilde{F}(\tilde{\xi})$ , we must add gravity term:

$$+ p_1 g$$

a. From Lesson #1, II.A.4.b.2, we have  $p_1 = \tilde{\xi} \cdot \nabla p_0 \equiv p_0 \nabla \cdot \tilde{\xi} = 0$

b. For incompressible motion,  $\nabla \cdot \tilde{\xi} = 0$ , so  $p_1 = -\tilde{\xi} \cdot \nabla p_0 = -\tilde{\xi}_x p_0'$

c. Thus  $+p_1 g = (-\tilde{\xi}_x p_0')(-g \hat{x}) = p_0' g \tilde{\xi}_x \hat{x}$

2. This gives:

$$\tilde{F}(\tilde{\xi}) = \nabla \left[ \tilde{\xi} \cdot \nabla p_0 + \gamma p_0 \nabla \cdot \tilde{\xi} \right] + \frac{(\tilde{\xi} \cdot \nabla p_0) \times [\nabla \times (\tilde{\xi} \times \tilde{B}_0)] + (\nabla \times [\nabla \times (\tilde{\xi} \times \tilde{B}_0)]) \times \tilde{B}_0}{\rho_0}$$
$$+ p_0' g \tilde{\xi}_x \hat{x}$$

3. For the energy principle, we must add this term:

$$-\frac{1}{2} |\tilde{\xi}_x|^2 g p_0'$$

$$[\text{Remember, } SW = -\frac{1}{2} \int \tilde{\xi}_x \cdot \tilde{\xi} \cdot \tilde{F}(\tilde{\xi})]$$

## Lecture #3 (Continued)

Hawes ⑥

### II. (Continued)

#### B. Using Energy Principle

1. With the added gravitational potential term, we have

$$SW = \frac{1}{2} \int d\mathbf{x} \left[ \frac{\nabla \times (\underline{\xi} + \underline{B}_0)^2}{\mu_0} + \gamma p_0 |\nabla \underline{\xi}|^2 \right] - \underline{\xi}^* \cdot \underline{j}_0 \times [\nabla \times (\underline{\xi} + \underline{B}_0)] \quad (1)$$

$$- \underline{\xi}^* \cdot \nabla (\underline{\xi} \cdot \nabla p_0) - |\underline{\xi}_x|^2 g p_0' \quad (2)$$

$$2. a. \nabla \times (\underline{\xi} + \underline{B}_0) = \underline{\xi} (\nabla \underline{B}_0) - \underline{B}_0 (\nabla \underline{\xi}) + (\underline{B}_0 \cdot \nabla) \underline{\xi} - (\underline{\xi} \cdot \nabla) \underline{B}_0 \quad (3)$$

NRL (10) p.4

$$= - \underline{B}_0' \underline{\xi}_x \hat{z}$$

$$b. \underline{j}_0 = \frac{\nabla \times \underline{B}_0}{\mu_0} = \frac{1}{\mu_0} (\partial_x \hat{x} + \partial_y \hat{y}) \times (\underline{B}_0 \hat{z}) = - \frac{1}{\mu_0} \underline{B}_0' \hat{y}$$

$$3. \text{TERM } (1) = \frac{(\underline{B}_0')^2}{\mu_0} |\underline{\xi}_x|^2$$

$$4. \text{TERM } (3) = \underline{\xi}^* \cdot \left( - \frac{\underline{B}_0'}{\mu_0} \hat{y} \right) \times (- \underline{B}_0' \underline{\xi}_x \hat{z}) = \underline{\xi}^* \cdot \left( \frac{(\underline{B}_0')^2}{\mu_0} \underline{\xi}_x \hat{y} \right) = - \frac{(\underline{B}_0')^2}{\mu_0} |\underline{\xi}_x|^2$$

a. Thus, Term (1) + Term (3) = 0 ✓

$$5. \text{Term } (4): - \underline{\xi}^* \cdot \nabla [(\underline{\xi} \cdot \nabla) p_0]$$

a. NOTE: NRL

$$(7) \text{ p.4 } \nabla \cdot (f A) = f \nabla \cdot A + A \cdot \nabla f$$

$$\nabla \cdot (\underline{\xi}^* f) = f \nabla \cdot \underline{\xi}^* + \underline{\xi}^* \cdot \nabla f$$

$$b. \text{Thus } \int d\mathbf{x} \left\{ \cancel{\underline{\xi}^* \cdot \nabla} - \underline{\xi}^* \cdot \nabla [(\underline{\xi} \cdot \nabla) p_0] \right\} = \int d\mathbf{x} \nabla \cdot \left[ \underline{\xi}^* (\underline{\xi} \cdot \nabla) p_0 \right]$$

$$\text{By Divergence Thm} = \oint dS \cdot \underline{\xi}^* (\underline{\xi} \cdot \nabla) p_0 = \oint dS \cdot [\underline{\xi}^* \underline{\xi}_x p_0'] = 0$$

NRL (28) p.5

By B.C. is  $\underline{\xi}_x$  at boundary  $\Rightarrow = 0$ , a  
is zero!

Periodic in  $y$  &  $z$  sums to zero.

6. Term (5): Only term left:

$$SW = - \frac{1}{2} \int d\mathbf{x} |\underline{\xi}_x|^2 g p_0'$$

Lecture #3 (Continued)  
II. B. (Continued)

Pages ⑦

7. Thus, for  $\rho_0' > 0$  (density increasing with height),

$\delta W < 0 \Rightarrow \text{UNSTABLE!}$