

# Hanes ①

## Lecture #11: Kinetic Theory for Electromagnetic Waves in Unmagnetized Plasma

### I. Review of Kinetic Theory

#### A. The Boltzmann Equation (Plasma Kinetic Equation)

$$1. \frac{\partial f_s}{\partial t} + \underline{v} \cdot \nabla f_s + \underbrace{\frac{q_s}{m_s} (\underline{E} + \underline{v} \times \underline{B})}_{\text{Lorentz Force}} \cdot \frac{\partial \underline{v}}{\partial \underline{x}} = \left( \frac{\partial f_s}{\partial t} \right)_{\text{coll}}$$

Lorentz Force gives acceleration

where  $f_s(\underline{x}, \underline{v}, t)$  is the distribution function in 6-D phase space.

2a. Recall from PHYS 4731 Lecture #11, ratio of collisional to collective effects in a plasma,

$$\frac{\text{Collisional effects}}{\text{Collective effects}} \sim \frac{N_e}{\omega_{pe}} \sim \frac{1}{N_0} \ll 1$$

Number of particles in the Debye sphere,

b. Thus, for many plasmas, we can neglect the effect of collisions,

$$\left( \frac{\partial f}{\partial t} \right)_{\text{coll}} = 0$$

This, Boltzmann Equation  $\Rightarrow$  Vlasov Equation

#### B. Vlasov-Maxwell System of Equations

i. The starting point for our studies of kinetic theory is this system.

$$a. \text{Vlasov Equation} \quad \frac{\partial f_s}{\partial t} + \underline{v} \cdot \nabla f_s + \underbrace{\frac{q_s}{m_s} (\underline{E} + \underline{v} \times \underline{B})}_{\text{Lorentz Force}} \cdot \frac{\partial \underline{v}}{\partial \underline{x}} = 0 \quad \text{for } s=1, e$$

$$b. \text{Maxwell's Equations:} \quad \nabla \cdot \underline{E} = \frac{\rho_2}{\epsilon_0} \quad \text{Poisson}$$

$$\rho_2 = \int_S d^3 v q_s f_s$$

$$\nabla \times \underline{E} = - \frac{\partial \underline{B}}{\partial t} \quad \text{Faraday}$$

$$\dot{B} = \int_S d^3 v q_s \underline{v} \times \underline{f}_s$$

$$\nabla \times \underline{B} = \mu_0 \underline{j} + \mu_0 \epsilon_0 \frac{\partial \underline{E}}{\partial t} \quad \text{Ampere-Maxwell}$$

$$\nabla \cdot \underline{B} = 0$$

c. This is a closed, integro-differential system of equations for 8 unknowns  $f_s(\underline{x}, \underline{v}, t)$ ,  $\underline{f}_e(\underline{x}, \underline{v}, t)$ ,  $\underline{E}(\underline{x}, t)$ ,  $\underline{B}(\underline{x}, t)$

## Lecture 11 (Continued)

Hanes 2

### I. (Continued)

#### C. The Distribution Function

##### 1. Maxwellian Distributions

a. for many linear problems, we take the bivariate order (equilibrium) distribution function to be Maxwellian.

b. Maxwellian distributions characterize Local Thermodynamic Equilibrium (a maximum entropy state — no free energy)

c.

$$f_{sm}(x, v, t) = \frac{n_s(x, t)}{\pi^{3/2} V_s^3} e^{-\frac{m_s(v - U(x, t))^2}{2 T_s(x, t)}}$$

where Def: Thermal Velocity  $V_s^2 = \frac{2 T_s(x, t)}{m_s}$

NOTE: Note that, for the rest of the semester, we will absorb Boltzmann's constant  $k = 1.38 \times 10^{-23} \frac{J}{K}$  into the temperature,  $kT \Rightarrow T$ , giving temperature  $T$  in units of energy (J).

d. For steady uniform conditions (homogeneous in space,  $\frac{\partial f}{\partial x} = 0$ ) and no flow velocity  $U = 0$ , this simplifies to

$$f_{sm}(v) = \frac{n_{so}}{\pi^{3/2} V_s^3} e^{-\frac{v^2}{V_s^2}}$$

NOTE: This is a function of only  $v = |v|$ .

##### 2. Moments of the distribution function:


a. Density:  $n_{so} = \int d^3 v f_{sm}(v)$

b. Energy:  $\frac{3}{2} n_{so} T_{so} = \int d^3 v \frac{1}{2} m_s v^2 f_{sm}(v)$


Lecture #11 (Continued)

T.C. (Continued)

3. Plot



Normalized by species thermal velocity



Equal areas under curves

For  $n_{i_0} = n_{e_0}$ .

4. Reduced Distribution Function:

a. DEF:  $F_s(v_z) \equiv \frac{1}{n_{s0}} \int_{-\infty}^{\infty} dv_x \int_{-\infty}^{\infty} dv_y f_{sm}(v)$

b. Integrate over two velocity space dimensions  $v_x$  &  $v_y$ .

c. For a Maxwellian,

$$F_s(v_z) = \frac{1}{n_{s0}} \int_{-\infty}^{\infty} d^3v \left[ \frac{n_{s0}}{\pi^{3/2} V_{Ts}^{3/2}} e^{-\frac{(v_x^2 + v_y^2 + v_z^2)}{V_{Ts}^2}} \right]$$

$$= \frac{e^{-\frac{v_z^2}{V_{Ts}^2}}}{\pi^{3/2} V_{Ts}} \underbrace{\int_{-\infty}^{\infty} dv_x}_{=1} \underbrace{\frac{e^{-\frac{v_x^2}{V_{Ts}^2}}}{\pi^{1/2}}}_{=1} \underbrace{\int_{-\infty}^{\infty} dv_y}_{=1} \underbrace{\frac{e^{-\frac{v_y^2}{V_{Ts}^2}}}{\pi^{1/2}}}_{=1} = \frac{e^{-\frac{v_z^2}{V_{Ts}^2}}}{\pi^{3/2} V_{Ts}} = F_s(v_z)$$

d. NOTE:  $\int_{-\infty}^{\infty} F_s(v_z) dv_z = 1$

## I. Electromagnetic Waves in an Unmagnetized Plasma

A. Setup

1. Electrostatic Approximation,  $B_1 = 0 \Rightarrow \nabla \times \vec{E} = 0 \Rightarrow \vec{E} = -\nabla \phi$

2. No mean magnetic field,  $B_0 = 0$ , And  $E_0 = 0$ .

3. Vlasov-Maxwell Systems Simplifies to

a.  $\frac{\partial f_s}{\partial t} + \vec{v} \cdot \nabla f_s - \frac{q_s}{m_s} \nabla \phi \cdot \frac{\partial \vec{r}}{\partial \vec{v}} = 0$  (for ions & electrons.)

b.  $-\nabla^2 \phi = \frac{1}{\epsilon_0} \leq \int d^3v q_s f_s$

Closed system of Equations for  $f_e(x, v, t)$ ,  $f_i(x, v, t)$ ,  $\phi(x, t)$

## II (Continued)

Hawes ④

### B. Linearization:

1. NOTE: Now  $\mathbf{v}$  is a coordinate and not a variable.

Therefore, we don't expand  $\mathbf{v}$ .

2. Take

$$a. f_s = f_{s_0}(\mathbf{v}) + \epsilon f_{s_1}(\mathbf{x}, \mathbf{v}, t)$$

$$b. \phi(\mathbf{x}, t) = \phi_0 + \epsilon \phi_1(\mathbf{x}, t)$$

3. NOTE: a. In steady state,  $\frac{\partial f_{s_0}}{\partial t} = 0$

b. For a homogeneous plasma,  $\frac{\partial f_{s_0}}{\partial \mathbf{v}} = 0$ .

c. Since  $E_0 = 0 \Rightarrow \phi_0 = \text{constant} \rightarrow 0$

$$4. a. \epsilon \frac{\partial f_{s_1}}{\partial t} + \epsilon \mathbf{v} \cdot \nabla f_{s_1} - \epsilon \frac{q_s}{m_s} \nabla \phi_1 \cdot \frac{\partial f_{s_0}}{\partial \mathbf{v}} - \epsilon^2 \frac{q_s}{m_s} \nabla \phi_1 + \frac{\partial f_{s_1}}{\partial \mathbf{v}} = 0$$

$$b. -\epsilon \nabla^2 \phi_1 = \frac{1}{\epsilon_0 s} \leq S d^3 v q_s f_{s_0} + \frac{1}{\epsilon_0 s} \leq S d^3 v q_s f_{s_1}$$

5. NOTE:  $S d^3 v q_s f_{s_0} = n_{s_0} q_s$ , so the first term of RHS of Poisson's equations becomes  $\frac{1}{\epsilon_0 s} \leq n_{s_0} q_s = 0$  for charge neutral equilibrium.

6. O( $\epsilon$ ):

$$\frac{\partial f_{s_1}}{\partial t} + \mathbf{v} \cdot \nabla f_{s_1} - \frac{q_s}{m_s} \nabla \phi_1 \cdot \frac{\partial f_{s_0}}{\partial \mathbf{v}} = 0$$

$$-\nabla^2 \phi_1 = \frac{1}{\epsilon_0 s} \leq S d^3 v q_s f_{s_1}$$

### C Fourier Transform in Space and Time

1. As usual, we'll solve this by Fourier transform  $\sim e^{i(k_x x - \omega t)}$

$$\nabla \Rightarrow i \underline{k} \quad \frac{\partial}{\partial t} \Rightarrow -i \omega$$

$$2. -i \omega f_{s_1} + i \mathbf{v} \cdot \underline{k} f_{s_1} - i \frac{q_s \phi_1}{m_s} \underline{k} \cdot \frac{\partial f_{s_0}}{\partial \mathbf{v}} = 0$$

$$k^2 \phi_1 = \frac{1}{\epsilon_0 s} \leq S d^3 v q_s f_{s_1}$$

## II.C (Continued)

Hawes (5)

3. Solving for  $f_{S1}$ :

$$f_{S1} = \frac{-q_s \phi_1}{m_s} \frac{k \cdot \frac{\partial f_{S0}}{\partial x}}{\omega - k \cdot v}$$

4. Substituting  $f_{S1}$  into Poisson's Equation:

$$a. k^2 \phi_1 = \frac{1}{\epsilon_0} \sum_s S d v_z \left( \frac{-q_s^2 \phi_1}{m_s} - \frac{k \cdot \frac{\partial f_{S0}}{\partial x}}{\omega - k \cdot v} \right)$$

5. Dividing by  $k^2$  and collecting terms:

$$\underbrace{\left[ 1 + \sum_s \left( \frac{n_s q_s^2}{\epsilon_0 m_s} \right) \frac{1}{k^2 n_s} \sum_s S d v_z \frac{k \cdot \frac{\partial f_{S0}}{\partial x}}{\omega - k \cdot v} \right]}_{\text{Dispersion relation } D(\omega, k)} \phi_1 = 0$$

Dispersion relation  $D(\omega, k)$

6. Simplifying

i. Take  $k = k \hat{z}$  without loss of generality.

$$a. k \cdot \frac{\partial f_{S0}}{\partial x} = k \frac{\partial f_{S0}}{\partial v_z}$$

$$b. k \cdot v = k v_z$$

$$c. D(\omega, k) = \left\{ 1 + \sum_s \frac{\omega_{ps}^2}{k^2} \int_{-\infty}^{\infty} dv_z \frac{k \frac{\partial}{\partial v_z} \left[ \int_{-\infty}^{\infty} dv_x \int_{-\infty}^{\infty} dv_y f_{S0} \right]}{\omega - k v_z} \right\} = 0$$

$= F_{S0}(v_z)$  Reduced Distribution Function

$$b. D(\omega, k) = 1 - \sum_s \frac{\omega_{ps}^2}{k^2} \int_{-\infty}^{\infty} dv_z \frac{\frac{\partial f_{S0}}{\partial v_z}}{v_z - \frac{\omega}{k}} = 0$$

Dispersion Relation

E. Failure of Fourier Transform Approach

i. The integral above in  $D(\omega, k)$  does not converge.

a. At  $v_z = \frac{\omega}{k}$ , denominator is zero.

b. Unless  $\frac{\partial f_{S0}}{\partial v_z} (and F_{S0}) = 0$  at  $v_z = \frac{\omega}{k}$ , integral does not converge.

## II. E. (Continued)

Haves ⑥

2. The failure at  $v_z = \frac{\omega}{k}$  occurs when there are particles with velocities that match the phase velocity of the wave.

⇒ These particles are resonant with the wave.

## F. The Bohm-Gross Dispersion Relation

1. Let's consider a plasma of electrons with stationary ions forming a neutralizing background.  $N_{i_0} = N_{e_0}$ ,  $f_i = 0$ .

### 2. Cold Plasma Limit:

- a. Assume electron thermal velocity much less than phase velocity.

$$v_{te}^2 \approx \langle v_z^2 \rangle \ll \frac{\omega^2}{k^2}$$

$$\text{where } \langle v_z^2 \rangle = \int_{-\infty}^{\infty} dv_z v_z^2 F_{e_0}(v_z)$$

- b. Strictly, we can only use this approach when  $F_{e_0}(v_z) \Big|_{v_z=\frac{\omega}{k}} = 0$   
(no resonant particles).

### 3a. Integrate $D(\omega, k)$ by parts

$$\int_{-\infty}^{\infty} dv_z \frac{\partial F_{e_0}}{\partial v_z} \frac{1}{v_z - \frac{\omega}{k}} = \left[ \frac{F_{e_0}}{v_z - \frac{\omega}{k}} \right]_{-\infty}^{\infty} + \int_{-\infty}^{\infty} dv_z \frac{F_{e_0}}{(v_z - \frac{\omega}{k})^2}$$

$$U = \frac{1}{v_z - \frac{\omega}{k}}$$

$$dU = \frac{\partial F_{e_0}}{\partial v_z} dv_z$$

$$dU = \frac{-dv_z}{(v_z - \frac{\omega}{k})^2}$$

$$V = F_{e_0}$$

$$\lim_{v_z \rightarrow \pm\infty} F_{e_0}(v_z) = 0$$

$$\text{b. Thus } D(\omega, k) = 1 - \frac{\alpha_{pe}^2}{k^2} \int_{-\infty}^{\infty} dv_z \frac{F_{e_0}}{(v_z - \frac{\omega}{k})^2} = 0$$

### 4. Expand denominator for $v_z \ll \frac{\omega}{k}$

$$\frac{1}{(v_z - \frac{\omega}{k})^2} = \frac{k^2}{\omega^2 (1 - \frac{k v_z}{\omega})^2} \approx \frac{k^2}{\omega^2} \left[ 1 + 2 \left( \frac{k v_z}{\omega} \right) + 3 \left( \frac{k v_z}{\omega} \right)^2 + \dots \right]$$

## II. F. (Continued)

Horwitz (?)

$$5. D(\omega, k) = 1 - \frac{\omega_{pe}^2}{\omega^2} \int_{-\infty}^{\infty} dv_z F_0 \left[ 1 + 2 \left( \frac{k v_z}{\omega} \right) + 3 \left( \frac{k v_z}{\omega} \right)^2 \right]$$

①      ②      ③

$$\text{①} = \int_{-\infty}^{\infty} dv_z F_0 = 1$$

$$\text{②} = \int_{-\infty}^{\infty} dv_z \frac{(k)}{\omega} V_z F_0 = 0 \quad (\text{odd in } V_z)$$

$$\text{C. ③} = 3 \frac{k^2}{\omega^2} \int_{-\infty}^{\infty} dv_z V_z^2 F_0(V_z) = \frac{3k^2}{\omega^2} \langle V_z^2 \rangle$$

NOTE: The form of  $\langle V_z^2 \rangle$  depends on equilibrium distribution function.

$$6. a. D(\omega, k) = 1 - \frac{\omega_{pe}^2}{\omega^2} \left( 1 + \frac{3k^2 \langle V_z^2 \rangle}{\omega^2} \right) = 0$$

$$b. \boxed{\omega^2 = \omega_{pe}^2 \left( 1 + 3 \frac{k^2 \langle V_z^2 \rangle}{\omega^2} \right)}$$

c. NOTE: We have assumed  $\frac{k^2 \langle V_z^2 \rangle}{\omega^2} \ll 1$ , so second term is a small correction. This can be solved easily by the method of successive approximations.

### 7. Method of Successive Approximations:

a. Solve for  $\omega^2$  by dropping small term:  $\omega^2 = \omega_{pe}^2 \left( 1 + \frac{3k^2 \langle V_z^2 \rangle}{\omega^2} \right)$   
 $\Rightarrow \omega_0^2 = \omega_{pe}^2$

b. Insert first solution  $\omega_0^2$  into small term to get second solution  $\omega_1^2$ :

$$\omega_1^2 = \omega_{pe}^2 \left( 1 + \frac{3k^2 \langle V_z^2 \rangle}{\omega_0^2} \right) = \omega_{pe}^2 + 3k^2 \langle V_z^2 \rangle$$

$\Rightarrow = \omega_{pe}^2$

c. Thus, we find  $\boxed{\omega^2 = \omega_{pe}^2 + 3k^2 \langle V_z^2 \rangle}$

## I. F. (Continued)

Haves ⑧

### 8. Alternative Explanation of Method of Successive Approximations

a.  $\omega^2 = \omega_{pe}^2 \left( 1 + \frac{3k^2 \langle v_z^2 \rangle}{\omega^2} \right)$

small term

b. Let  $x = \omega^2$ ,  $a = \omega_{pe}^2$ ,  $b = 3k^2 \langle v_z^2 \rangle \Rightarrow x = a(1 + \frac{b}{x})$

c.  $x = x_0 + \epsilon x_1 + \epsilon^2 x_2 + \dots$

d.  $x_0 + \epsilon x_1 = a \left[ 1 + \epsilon \frac{b}{x_0 + \epsilon x_1} \right] = a \left[ 1 + \frac{\epsilon b}{x_0(1 + \epsilon x_1)} \right] \approx a \left( 1 + \frac{\epsilon b}{x_0} - \epsilon^2 \frac{bx_1}{x_0^2} \right)$

e.  $O(1)$ :  $x_0 = a \Rightarrow \omega_0^2 = \omega_{pe}^2$

f.  $O(\epsilon)$ :  $x_1 = \frac{ab}{x_0} \Rightarrow \omega_1^2 = \frac{\omega_{pe}^2 3k^2 \langle v_z^2 \rangle}{x_0^2} = 3k^2 \langle v_z^2 \rangle$

g.  $x = x_0 + \epsilon x_1 \Rightarrow \omega^2 = \omega_{pe}^2 + 3k^2 \langle v_z^2 \rangle$

### 9. Maxwell Equilibrium Distribution:

$$\langle v_z^2 \rangle = \int_{-\infty}^{\infty} dv_z v_z^2 F_{e0}(v_z) = \int_{-\infty}^{\infty} \frac{dv_z}{V_{Te}} \frac{v_z^2}{V_{Te}^2} e^{-\frac{v_z^2}{V_{Te}^2}} = V_{Te}^2 \int_0^{\infty} dy y^2 e^{-y^2}$$

$$= V_{Te}^2 \frac{\sqrt{\pi}}{2} = \left( \frac{8T_e}{m_e} \right)^{\frac{1}{2}} = \frac{T_e}{m_e} \quad Y = \frac{v_z}{V_{Te}}$$

b. Thus  $\boxed{\omega^2 = \omega_{pe}^2 + 3k^2 \frac{T_e}{m_e}}$  Langmuir Waves

c. From PHYS 4731 Lec #24,  $\omega^2 = \omega_{pe}^2 + k^2 C_e^2$  for Langmuir Waves

1. DEF: Electron Sound Speed:  $C_e^2 = \frac{T_e}{m_e}$

2. In Lecture #24, we took  $\gamma_e = 1$  for isothermal conditions, which requires  $V_{Te} \gg \frac{k}{\lambda}$ . Here we are in the opposite limit, so we put  $\gamma_e$  back in.

### 10. Bohm-Gross Dispersion Relation

a.  $\omega^2 = \omega_{pe}^2 + \gamma_e k^2 C_e^2$  Warm, Two-Fluid Theory (Fluid Conservation)

b.  $\omega^2 = \omega_{pe}^2 + 3k^2 C_e^2$  Kinetic Theory

c. Results agree for  $\gamma_e = 3$  (one degree of freedom,  $\gamma = \frac{F+2}{F}$ )

AAA Kinetic Theory gives result without assuming an Equation of State!