### Review: Newton's 1st & 2nd Laws

- 1st law (Galileo's principle of inertia)- no force is needed to keep an object moving with constant velocity
- 2<sup>nd</sup> law (law of dynamics) a net force must be applied to change the velocity of an object.
- A force F (N) = m (kg) x a (m/s²) must be applied to produce an acceleration a for an object of mass m

# L-7 Newton's third law and conservation of momentum

For every action there is an equal and opposite reaction.

For Newton: Action = Force

.

### Forces: pushes or pulls

- If I push against a person, that person pushes back on me
- If I push on the wall, the wall pushes back on me, otherwise I fall over!
- You can't push on something unless it pushes back on you!
- Forces never occur alone, they occur in pairs
- Every force is part of an interaction


# A boxer's punch




- The boxer's fist hits the bag and the bag hits back on the fist → a pair of BIG forces are involved in hitting the bag
- When the boxer hits a piece of tissue paper, the boxer can only exert as much force on the tissue paper as the tissue paper can exert on his fist

## Newton's 3rd Law

Whenever one object (A) exerts a force on a second object (B), the second object exerts an equal and opposite force on the first.




# Example: static equilibrium



- What keeps the box on the table?
- The box exerts a force on the table due to its weight, and as result of the 3<sup>rd</sup> law the table exerts an equal and opposite (upward) force on the box.
- If the table was not strong enough to support the weight of the box, the box would crash through it.

## Example: The bouncing ball

- · Why does the ball bounce?
- · When the ball hits the ground it exerts a downward force on it
- By the 3rd law, the ground must exert an equal and upward force on the ball
- The ball bounces because of the upward force exerted on it by the ground.



# • Since the earth exerts a downward

You can move the earth!

- force on you, the 3<sup>rd</sup> law requires that you exert an equal upward force on the earth.
- Because the earth's mass and your mass are different, the acceleration of the earth and your acceleration are NOT equal
- The Earth's acceleration is much less than your acceleration because the mass of the earth is much bigger than your mass.



- Newton's 3<sup>rd</sup> Law plays an important role in everyday life, whether we realize it or not.
- · We will demonstrate this in the next few examples.


# The donkey and 3rd law paradox

· A man tries to make a donkey pull a cart but the donkey argues:



- Why should I even try?
- No matter how hard I pull on the cart, the cart pulls back on me with an equal force, so I can never move it. What is the fallacy in the donkey's argument?
- The donkey forgot that action/reaction forces always act on different objects. As far as the cart is concerned, if the force the donkey exerts on it is large enough, it will move. The reaction force on him is irrelevant.

#### Friction is essential to movement




- The tires push back on the road and the road pushes the tires forward.
- If the road is slippery, the friction force between the tires and road is reduced, and the car does not move.

## We could not walk without friction




- · When we walk, we push back on the ground
- By the 3<sup>rd</sup> law the ground then pushes us forward.
- If the ground is slippery, we cannot push back on it, so it cannot push forward on us  $\rightarrow$  we go nowhere!




## Bouncing and Non-bouncing balls

- The ball that bounced exerted the larger force on the ground.
- The force that the ball exerts on the ground is equal to and in the opposite direction as the force of the ground on the ball.
- The ground must exert a force to bring the non-bouncy ball to rest
- The ground must not only stop the bouncy ball, but must then project it back up → this requires more force!
- Since the bouncy ball experiences a larger force from the ground, it must therefore by the 3<sup>rd</sup> Law also exert a larger force ON the ground.
- The next demo should convince you!

14





#### But why does the beaker break?

- An object will break if a large enough force is exerted on it.
   Obviously, the beaker that hits the brick experiences the larger force, but can we explain this using Newton's Laws?
- Notice that in both cases, the beakers have the same velocity
  just before hitting the cushion or the brick. Also both beakers
  come to rest (one gently, the other violently) so their final
  velocities are both zero.
- Both beakers therefore experience the same *change in* velocity =  $\Delta v$  (delta  $\Delta$  means change), so that the change in v  $\Delta v = v_f v_i = 0 v_i = -v_i$
- What about their acceleration?
- a = Δv / Δt, where Δt is the time interval over which the velocity changes (the time to stop)



#### Continued from previous slide

- The stopping time  $\Delta t$  is the important parameter here because it is *not the same* in both cases.
- The beaker falling on the cushion takes longer to stop than the one falling on the brick. The cushion allows the beaker to stop gently, while the brick stops it abruptly
- Both beakers have the same velocity just before hitting the bottom, and both come to rest, so the change in velocity is the same in both cases.
- However, the beaker that hits the brick, is stopped suddenly and thus experiences a greater acceleration and a greater force which cause it to shatter.

18

# Stunt actors and air bags

- The same reasoning applies to stunt actors and air bags in an automobile. <u>Video</u>
- Air bags deploy very quickly, triggered when an unusually large acceleration is detected
- They provide protection by allowing you to stop more slowly, as compared to the case where you hit the steering wheel or the windshield.
- Since you come to rest more slowly, the force on you is smaller than if you hit a hard surface.
- Some say: "Airbags slow down the force."
   Although this is not technically accurate, it is a good way of thinking about it.

#### Momentum

- Newton's 3<sup>rd</sup> law leads directly to the concept of (linear) momentum
- Momentum is a term used in everyday conversation, e.g., "The team has momentum," or, "The team lost its momentum."
- These phrases imply that if you get momentum, you tend to keep it, but when you loose it, it is hard to get it back.
- This colloquial usage is similar to a concept we will discuss – the law of conservation of momentum – a law that is very useful in understanding what happens when objects collide.

20


# Physics definition of momentum (p)

- In physics, every quantity must be unambiguously defined, with a prescribed method for measurement
- Momentum (p) = mass (m) x velocity (v)
- Based on the simple observation that a heavy truck is harder to stop than a small car at the same speed
- p = m v (m in kg, v in m/s, p in kg m/s)
- An object with small m and large v, can have a comparable momentum as a very massive moving more slowly.
- e. g., (a) m = 2 kg, v = 10 m/s → p = 20 kg m/s
   (b) m = 5 kg, v = 4 m/s → p = 20 kg m/s

21

#### Conservation of momentum in collisions

- The collision of 2 objects can be very complicated; large forces are involved (which usually cannot be measured) which act over very short time intervals
- Even though the collision forces are not known, the 3<sup>rd</sup> law ensures that the forces that the objects exert on each other are equal and opposite – this is a very big simplification
- Application of Newton's 3<sup>rd</sup> and 2<sup>nd</sup> laws leads to the conclusion that in the collision, the total momentum of the two objects before the collision is the same as their total momentum after the collision – this is called conservation of momentum.



- The sum of the momentum of A and B before the collision = the sum of the momentum of A and B after the collision.
- In the next lecture, we will discuss other types of collisions

23