

The light-front vacuum and dynamics*

W. N. Polyzou

The University of Iowa, Iowa City, IA

Abstract. I give a quantum theoretical description of kinematically invariant vacua on the algebra of free fields restricted to a light front and discuss the relation between the light-front Hamiltonian, P^- , the vacuum, and Poincaré invariance. This provides a quantum theoretical description of zero modes.

1 Introduction

A light-front field, $\phi(f)$, is a free field restricted to the light front $x^+ = 0$ and smeared with a Schwartz test function of the light-front variables $\tilde{x} = (x^-, \mathbf{x}_\perp)$. The commutator of two light-front fields is:

$$[\phi(f), \phi(g)]_- = \frac{1}{2}[(f, g)_f - (g, f)_f] \quad (f, g)_f := \int \frac{d\tilde{p}\theta(p^+)}{p^+} \tilde{f}^*(\tilde{p})\tilde{g}(\tilde{p}) \quad (1)$$

where $\tilde{f}(\tilde{p})$ is the Fourier transform of $f(\tilde{x})$. In the absence of restrictions on the test functions, the light-front scalar product in (1) diverges logarithmically due to the $p^+ = 0$ singularity in the denominator. The commutator (1) becomes finite if the test functions are restricted to have the form $\tilde{f}(\tilde{p}) = p^+ \tilde{g}(\tilde{p})$, where $\tilde{g}(\tilde{p})$ are ordinary Schwartz test functions of the light-front momenta. This space of test functions, introduced by Schlieder and Seiler [1], is denoted by \mathcal{S}^+ .

The light-front Fock algebra, \mathcal{A}_f , is generated by finite linear combinations of the form

$$A := \sum_{k=1}^N c_k e^{i\phi(f_k)} \quad (2)$$

where c_k are complex and $f_k(\tilde{x})$ are real Schlieder-Seiler functions. It is straightforward to show that \mathcal{A}_f is an abstract $*$ -algebra with following properties:

1. \mathcal{A}_f is closed under kinematic Poincaré transformations.
2. \mathcal{A}_f is a Weyl algebra.

*Presented at Light-Cone 2004, Amsterdam, 16 - 20 August

3. Rotations induce algebraic isomorphisms from \mathcal{A}_f to algebras associated with different light-front orientations.

A vacuum is a positive, invariant, linear functional $E[\cdot]$ on this algebra. The properties of a vacuum can be expressed in terms of its generating functional

$$S\{f\} := E[e^{i\phi(f)}] = \langle 0|e^{i\phi(f)}|0\rangle.$$

The generating functional of a light-front vacuum must be normalized, $S\{0\} = 1$, real $S^*\{f\} = S\{-f^*\}$, continuous

$$f_n \rightarrow f \in \mathcal{S}^+ \Rightarrow S\{f_n\} \rightarrow S\{f\},$$

non-negative

$$S\{f_i - f_j\} := M_{ij} \geq 0$$

for any sequence $\{f_n\}$ of real test functions in \mathcal{S}^+ , kinematically invariant

$$S\{f\} = S\{f'\}$$

where $f'(x) = f(\Lambda x + a)$ for any *kinematic* Poincaré transformation, and satisfy cluster properties

$$\lim_{\lambda \rightarrow \infty} S\{f + g_\lambda\} \rightarrow S\{f\}S\{g\}$$

where $g_\lambda(x) = g(x + \lambda y)$ and y is any space-like vector in the light-front hyperplane [2].

The Hilbert space representation of \mathcal{A}_f associated with a given vacuum functional is defined as follows. A dense set of vectors is given by expressions of the form

$$|\psi\rangle = \sum_{n=1}^N c_n e^{i\phi(f_n)} |0\rangle \quad N < \infty.$$

The inner product can be expressed in terms of the generating functional

$$\langle \xi | \psi \rangle := \sum_{mn} d_m^* c_n e^{\frac{1}{2}[\phi(g_m), \phi(f_n)]} S\{f_n - g_m\}.$$

The generating functional of the Fock representation, $S_0\{f\} := e^{-\frac{1}{4}(f,f)}$, satisfies all of the required properties. The representation of the kinematic Poincaré transformations on this Hilbert space is unitary.

The algebra \mathcal{A}_f has another class of kinematically invariant vacua. Given a Schlieder-Seiler test function $\tilde{f}(\tilde{p})$ define

$$\hat{f}(\mathbf{p}_\perp) = \lim_{p^+ \rightarrow 0} \frac{\tilde{f}(p^+, \mathbf{p}_\perp)}{p^+}. \quad (3)$$

Vacuum generating functionals have the form

$$S\{f\} = S_0\{f\} s\{\hat{f}\} \quad s\{\hat{f}\} = e^{\sum_n i^n s_n(\hat{f}, \dots, \hat{f})} \quad (4)$$

where

$$s_n(\mathbf{p}_{1\perp}, \dots, \mathbf{p}_{n\perp}) := \delta\left(\sum_{i=1}^n \mathbf{p}_{i\perp}\right) w_{tn}(\mathbf{p}_{1\perp}, \dots, \mathbf{p}_{n\perp}) \quad (5)$$

and $w_{tn}(\mathbf{p}_{1\perp}, \dots, \mathbf{p}_{n\perp})$ are connected, two-dimensional, Euclidean invariant Schwartz distribution in $2(n-1)$ independent variables. The functional, $s\{\hat{f}\}$, is the Fourier transform of a positive measure on the cylinder sets of Schwartz distributions in two variables [3]. The generating function $S\{f\}$ will be the generating function of a vacuum functional if $s\{\hat{f}\}$ satisfies the same properties as $S\{f\}$ with respect to Schwarz functions of two variables, where the invariant subgroup is the two-dimensional Euclidean group. While positivity is a strong constraint, non-trivial examples associated with coherent states and Gaussian measures exist. The different vacuum generating functionals lead to inequivalent Hilbert space representations of the $*$ algebra.

To construct a relativistic model it is necessary to complete the Lie algebra of the Poincaré group by finding dynamical generators that are compatible with a given vacuum state. The first step is to construct a mass operator that is compatible with the vacuum. Let O be any non-negative kinematically invariant operator. The Fock representation mass operator, M_f , is one such example; others can be constructed by choosing $O = B^\dagger B$ for a kinematically invariant operator B . Let $\Pi := (I - |0\rangle\langle 0|)$.

The operator $M := \Pi O \Pi$ is non-negative, kinematically invariant, and annihilates the vacuum. It is a suitable candidate for the mass operator of a unitary representation of the Poincaré group. The associated light-front Hamiltonian P^- is:

$$P^- := \frac{\mathbf{P}_\perp \mathbf{P}_\perp + M^2}{P^+}.$$

The mass operator M can be formally expressed as a limit of elements of \mathcal{A}_f . To do this let $\{A_n\} \in \mathcal{A}_f$ generate an orthonormal basis for the Hilbert space representation with vacuum functional E , $E[A_n^\dagger A_m] = \delta_{mn}$. It follows that

$$M = \sum_{k,l} A_k^\dagger A_l m_{kl} \quad \text{where}$$

$$m_{kl} := E[A_k^\dagger O A_l] - E[A_k^\dagger] E[O A_l] - E[A_k^\dagger O] E[A_l] + E[A_k^\dagger] E[O] E[A_l]$$

which explicitly exhibits M as the limit of elements of the algebra.

The last step in constructing a dynamics is to complete the Lie algebra of the Poincaré group by including rotations. The ability to complete the Lie algebra is intrinsic to the choice of kinematically invariant M . Free rotations, $U_0(R)$, act on the fields covariantly and define algebraic isomorphisms from \mathcal{A}_f to light-front Fock Algebras with *different* light fronts. Given M and the kinematic observables it is possible to formulate a scattering theory. The point eigenstates of the mass operator are needed to formulate the scattering asymptotic condition. They are acceptable if under *free* rotations they satisfy

$$U_0(R)|m\rangle_{\hat{n}} = |m\rangle_{R\hat{n}} D(R)$$

where $D(R)$ is an irreducible representation of the rotation group (note the change in the orientation of the light front). Wave operators, constructed using these one-body solutions to formulate the asymptotic condition, satisfy

$$U_0(R)\Omega_{\hat{n}\pm} = \Omega_{R\hat{n}\pm} U_f(R)$$

where $\Omega_{R\hat{n}\pm}$ is the wave operator associated with the rotated light front, $R\hat{n}$ and $U_f(R)$ is the asymptotic representation of $SU(2)$. Poincaré invariance requires rotation operators that leave the vacuum invariant. If the scattering operators are asymptotically complete and *independent* of the orientation of the light-front,

$$\Omega_{+\hat{n}}^\dagger \Omega_{-\hat{n}} = \Omega_{+R\hat{n}}^\dagger \Omega_{-R\hat{n}} \Rightarrow A_R := \Omega_{+R\hat{n}} \Omega_{+\hat{n}}^\dagger = \Omega_{-R\hat{n}} \Omega_{-\hat{n}}^\dagger$$

then it follows that

$$U_{\hat{n}}(R) := U_0(R) A_{R^{-1}} = U_0(R) \Omega_{R^{-1}\hat{n}\pm} \Omega_{\hat{n}\pm}^\dagger = \Omega_{\hat{n}\pm} U_f(R) \Omega_{\hat{n}\pm}^\dagger$$

extends the Poincaré group on the Hilbert space with the given non-trivial vacuum [4]. The invariance of the S matrix ensures that A_R does not depend on the choice of asymptotic condition. Here wave operators are two-Hilbert space wave operators that necessarily include the point spectrum contributions to the mass operator. The \hat{n} -invariance of S can be tested in calculations, however the restrictions are non-trivial.

The above construction shows that the singularities in the light-front scalar product lead to a restriction on the test functions in the light-front Fock algebra. The resulting $*$ -algebra, \mathcal{A}_f , has a large class of kinematically invariant vacua that lead to inequivalent Hilbert space representations of the algebra. It is possible to find dynamical P^- or mass operators M that are positive, annihilate the vacuum and are kinematically covariant (resp. invariant). A large class of such operators exist, but a given operator can only annihilate one of the vacuum vectors. The dynamical operator P^- or M can be used to formulate a scattering theory. A sufficient condition for full Poincaré invariance is that the scattering matrix associated with different light fronts is independent the light front. This is a strong, but testable condition.

The construction provides a direct means to formulate dynamical models associated with inequivalent vacuum representations. This construction does not utilize classical equations of motion or models with a finite number of degrees of freedom, it provides a provide direct means to formulate and study models with non-trivial zero modes.

References

1. S. Schlieder and E. Seiler, Comm. Math. Phys **25**,62(1972)
2. F. Coester and W. N. Polyzou, Foundations of Physics **24**, 387(1994)
3. I. M. Gelfand and N. Ya. Vilekin, Generalized Functions, Vol. 4, Academic Press, (1964)
4. W. N. Polyzou, Few Body Systems, **27**,57(1999)