

Relativity and the low energy nd A_y puzzle

H. Witała, J. Golak, and R. Skibiński

*M. Smoluchowski Institute of Physics,
Jagiellonian University, PL-30059 Kraków, Poland*

W. Glöckle

Institut für theoretische Physik II, Ruhr-Universität Bochum, D-44780 Bochum, Germany

W. N. Polyzou

*Department of Physics and Astronomy,
The University of Iowa, Iowa City, IA 52242*

H. Kamada

*Department of Physics, Faculty of Engineering,
Kyushu Institute of Technology, Kitakyushu 804-8550, Japan*

(Dated: December 20, 2007)

Abstract

We solve the Faddeev equation in an exactly Poincaré invariant formulation of the three-nucleon problem. The dynamical input is a relativistic nucleon-nucleon (NN) interaction that is exactly on-shell equivalent to the high precision CD Bonn NN interaction. S -matrix cluster properties dictate how the two-body dynamics is embedded in the three-nucleon mass operator (rest Hamiltonian). We find that for neutron laboratory energies above ≈ 20 MeV relativistic effects on A_y are negligible. For energies below ≈ 20 MeV dynamical effects lower the nucleon analyzing power maximum slightly by $\approx 2\%$ and Wigner rotations lower it further up to $\approx 10\%$ increasing thus disagreement between data and theory. This indicates that three-nucleon forces (3NF) must provide an even larger increase of the A_y maximum than expected up to now.

PACS numbers: 21.45.+v, 24.70.+s, 25.10.+s, 25.40.Lw

I. INTRODUCTION

High precision nucleon-nucleon (NN) interactions such as AV18 [1], CDBonn [2], Nijm I, II and 93 [3] accurately describe the NN data set up to about 350 MeV. When these interactions are used to predict binding energies of three-nucleon (3N) systems they underestimate the experimental bindings of 3H and 3He by about 0.5-1 MeV [4, 5]. This missing binding energy can be cured by introducing a three-nucleon force (3NF) into the nuclear Hamiltonian [5].

The study of elastic nucleon-deuteron (Nd) scattering and nucleon-induced deuteron breakup also revealed a number of cases where the non-relativistic description based on pairwise interactions is insufficient to explain the data. Generally, the studied discrepancies between a theory based on NN interactions only and experiment become larger with increasing energy of the 3N system. Adding a 3NF that includes long-range 2π exchange to the pairwise interactions leads in some cases to a better description of the data. The parameters of such a 3NF must be separately adjusted to the experimental binding of 3H and 3He [6–8] for each NN interaction. The elastic Nd angular distribution in the region of its minimum and at backward angles is the best studied example [6, 8]. The clear discrepancy in these angular regions at energies below ≈ 100 MeV nucleon laboratory energy between a theory based on NN interactions only and the cross-section data can be removed by adding modern 3NFs based on chiral effective field theory [9] to the nuclear Hamiltonian. At energies higher than ≈ 100 MeV current 3NFs [10, 11] only partially improve the description of cross section data and the remaining discrepancies, which increase with energy, indicate the possibility of relativistic effects [12–14]. The need for a relativistic description of 3N scattering was also raised when precise measurements of the total cross section for neutron-deuteron (nd) scattering [15] were analyzed within the framework of non-relativistic Faddeev calculations [16]. NN interactions alone were insufficient to describe the data above ≈ 100 MeV.

In few-body models off-shell effects, relativistic effects, and three-body force contributions cannot be cleanly separated. This is because different two-body interactions that give the same two-body S matrix are related by a unitary scattering equivalence [17]. To maintain this equivalence at the three-body level requires additional three-body interactions [18] in one of the Hamiltonians. Since relativistic two-body models are fit to the same data as the corresponding non-relativistic models, there is a similar on-shell two-body scattering equivalence. While the relativistic and non-relativistic three-body predictions will be different one can in principle make up the difference (in a chosen frame) with a suitable three-body interaction. So while it is possible to simulate “relativistic effects” with a three-body interaction, a Poincaré invariant treatment of the dynamics provides the

most direct way to model the consequences of imposing Poincaré invariance and S-matrix cluster properties.

In this paper we investigate one particular representation of the Poincaré invariant three-body problem. We compare the predictions of relativistic and non-relativistic three-body calculations where the input two-body interactions give the same two-body S matrix, have the same internal two-body wave functions, and have a kinematic three-dimensional Euclidean symmetry. The relativistic and non-relativistic models differ in how these interactions appear in the three body-problem. In addition, the internal and single particle variables are related by Galilean boosts in the non-relativistic case and Lorentz boosts in the relativistic case. In the non-relativistic case the two-body interactions for each pair are added to the center of mass kinetic energy operator. In the relativistic case, the non-linear relation between the two and three-body mass operators must be respected in order to obtain a scattering matrix that clusters into a product of the identity and the input two-body S -matrix [19]. This non-linear dependence of the three-body invariant mass operator on the two-body interaction has dynamical consequences for the three-body system which complicates the structure of the Faddeev kernel.

In [20] we used a Poincaré invariant formulation of the 3N scattering problem. A technique for constructing the relativistic nucleon-nucleon interaction from a standard high-precision interaction was given in [21]. We used the same technique to construct the transition operators that appear in the kernel of the relativistic Faddeev equation. Application to a 3N bound state supported the relativistic effects previously found in [22].

Realistic NN interactions are fit by properly transforming experimental data to the center of momentum frame and fitting S -matrix elements computed using the non-relativistic Schrödinger equation to this data. While the same data could be precisely fit using S -matrix elements computed from a relativistic Schrödinger equation, this has not been done [23] with the same precision used to construct realistic interactions.

In [21], instead, an analytical scale transformation of momenta was used to relate NN interactions in the non-relativistic and relativistic Schrödinger equations in such a way, that the two-body scattering matrix elements are identified as $S_{nr}(E_{cm}) = S_r(E_{cm}) = S_{exp}(E_{cm})$ as functions of the center of momentum energy [24]. In this work we use an alternative procedure [25] that generates a relativistic nucleon-nucleon interaction with the property that the relativistic and non-relativistic two-body S matrices satisfy $S_r(\mathbf{k}^2) = S_{nr}(\mathbf{k}^2) = S_{exp}(\mathbf{k}^2)$, where $S_{exp}(\mathbf{k}^2)$ is the experimental two-body S matrix, and \mathbf{k}^2 is the cm momentum of one of the particles.

When high-precision potentials are determined [26] by properly Lorentz transforming scattering

data from the laboratory frame to the center of momentum frame the Lorentz invariant scalar product $p_{target} \cdot p_{beam} = mE_b = 2\mathbf{k}^2 + m^2$ is used to relate the laboratory beam energy E_b to the c.m. momentum \mathbf{k}^2 . The potential is determined by comparing the transformed experimental scattering observables to the scattering observables computed using the non-relativistic Lippmann-Schwinger equation, identifying the \mathbf{k}^2 appearing in the Lippmann-Schwinger equation with the \mathbf{k}^2 computed from the invariant $p_{target} \cdot p_{beam}$. With this procedure the resulting interactions are constructed so the S-matrix elements in the relativistic and non-relativistic cases are identified as functions of \mathbf{k}^2 rather than cm energy. Though the difference in the two approaches leads to a small mismatch in the relativistic and non-relativistic momentum, the interactions generated by the analytic scale transformation provide a useful first step to investigate the effects of the non-linear relation between the three-body mass operator and two-body interactions.

In our initial studies [20, 27, 28] the interaction generated by the analytic scale transformation was used to study the changes in elastic nd scattering and breakup observables when the non-relativistic form of the kinetic energy is replaced by the relativistic one and a proper treatment of the dynamics is included. We found that the elastic scattering cross section is only slightly influenced by relativity. Only at backward angles and higher energies the elastic cross sections are increased by relativity.

Due to the selectivity of the breakup reaction, however, regions of phase-space were found at higher energies of the incoming nucleon where relativity leads to a characteristic pattern by which relativity changes the non-relativistic breakup cross section. Namely, in this region of phase-space fixing the angle of the first detected nucleon and changing the angle of the second nucleon provides variations of the non-relativistic cross section by relativity, increasing or decreasing it by a factor up to ≈ 2 . For spin observables the implemented relativistic features lead only to small effects.

Recently an interesting estimate of “relativistic corrections” has been performed and its effect on low-energy nd analyzing power A_y has been estimated using the plane-wave impulse approximation [29]. The estimate is based on a perturbative realization of the Poincaré Lie algebra to leading order in $1/c^2$ [30]. The calculations are done in the plane-wave impulse approximation. A large increase by $\approx 10\%$ of the A_y maximum at laboratory energy $E_n = 3$ MeV has been found. The authors comment that their estimates are both exploratory and incomplete. In addition to the absence of final state interaction, there are a number of other important differences with an exact formulation of this problem. Such a large effect, which would significantly reduce the discrepancy between theory and data in the region of the A_y maximum, calls for a relativistic study in an exactly Poincaré invariant treatment of the three nucleon dynamics.

Our previous study [20], performed without inclusion of Wigner rotations, is too limited for spin observables. Therefore, in order to make definite conclusions for A_y we perform a complete dynamical calculation including the effects of Wigner rotations. We focus on that issue and do not include 3NF's.

The paper is organized as follows. In section II we discuss the construction of our Poincaré invariant dynamical model. This includes a discussion of how high-precision interactions are used to construct the three-body mass operator (rest Hamiltonian). In section III we discuss spin observables in Poincaré invariant quantum theory. In Sec. IV we discuss the formulation of the Faddeev equation to construct scattering observables for this three-nucleon mass operator. This includes an exact treatment of the Faddeev kernel which avoids the approximations used in [20, 27, 28]. We solve the relativistic 3N Faddeev equation with and without Wigner spin rotations. We show and discuss results for the neutron analyzing power A_y . Sec. V contains a summary and conclusions.

II. POINCARÉ INVARIANT DYNAMICS

In a quantum theory the principle of special relativity requires that the probabilities computed for equivalent experiments done in different inertial coordinate systems are identical. Since inertial coordinate systems are related by Poincaré transformations, it follows [31] that equivalent states in different inertial coordinate systems are related by a unitary representation, $U(\Lambda, a)$, of the Poincaré group. This emphasis on the invariance of experimental measurements in different inertial frames is different than the covariance requirements that are historically motivated by the way symmetries are realized in classical wave equations.

Since any representation of the Poincaré group can be decomposed into a direct integral of irreducible representations, one way to construct a Poincaré invariant dynamics is to build it out of irreducible representations. The transformation properties of irreducible representations are well known and completely determined by group theoretical considerations. The dynamics is contained in the spectrum of the physical mass and spin operators which determines the values and multiplicities of the Casimir invariants that appear in this decomposition.

Our construction begins with one-particle representations, which are irreducible representations. The particle's mass and spin fix the eigenvalues of the two Casimir invariants of the Poincaré group. For computations it is necessary to choose a basis for the irreducible representation space. This is done by choosing a maximal set of commuting Hermitian functions of the infinitesimal

Poincaré generators. In addition to the mass and spin, it is possible to find four additional mutually commuting non-invariant functions of the generators. There is a second set of four operators that are conjugate to the non-invariant commuting observables. These operators change the eigenvalues and determine the spectrum of the commuting observables. All ten generators can be expressed as functions of these eight non-invariant operators and the two Casimir operators. The irreducible representation space, \mathcal{H} , is the space of square integrable functions of the eigenvalues of the four commuting operators [32].

Our choice of basis for irreducible representation spaces is the simultaneous eigenstates of the linear momentum \mathbf{p} and the 3-component of the canonical spin, j_{cz} , which is the observable corresponding to the spin measured in the particle's rest frame if the particle is transformed to its rest frame with a rotationless Lorentz transformation. In this basis the irreducible unitary representation of the Poincaré group is [32]:

$$U(\Lambda, a)|j, m\rangle_{\mathbf{p}, \mu} = |(j, m)_{\mathbf{p}', \mu'}\rangle e^{ip' \cdot a} \sqrt{\frac{\omega(p')}{\omega(p)}} D_{\mu' \mu}^j [B^{-1}(\mathbf{p}'/m)\Lambda B(\mathbf{p}/m)] \quad (1)$$

where $p' = \Lambda p$, $\omega(p) = \sqrt{\mathbf{p} \cdot \mathbf{p} + m^2}$, and $B(\mathbf{p}/m)$ is the rotationless Lorentz transformation that takes a particle of mass m at rest to momentum \mathbf{p} . The quantity $R_w(\Lambda, p) := B^{-1}(\mathbf{p}'/m)\Lambda B(\mathbf{p}/m)$ is the standard rotationless-boost Wigner rotation. The representation (1) is unitary for states with a $\delta(\mathbf{p} - \mathbf{p}')$ normalization in the momentum. The important observation is that *all* mass $m > 0$ spin j irreducible representations of the Poincaré group in the $\{\mathbf{p}, j_{cz}\}$ basis have this form.

The two or three-nucleon Hilbert space is the tensor product of two or three single-nucleon irreducible representation spaces; $\mathcal{H} \otimes \mathcal{H}$ or $\mathcal{H} \otimes \mathcal{H} \otimes \mathcal{H}$. On each of these spaces

$$U_0(\Lambda, a) = U(\Lambda, a) \otimes U(\Lambda, a) \quad U_0(\Lambda, a) = U(\Lambda, a) \otimes U(\Lambda, a) \otimes U(\Lambda, a) \quad (2)$$

define kinematic representations of the Poincaré group. These representations are reducible and do not contain any dynamics. We build dynamical irreducible representations by adding suitable interactions to the mass Casimir operator of non-interacting irreducible representations. The first step needed to introduce interactions is then to decompose these non-interacting tensor product representations into a direct integral of irreducible representations. This is accomplished with Poincaré group Clebsch-Gordan coefficients in our chosen $\{\mathbf{p}, j_{cz}\}$ basis. The Poincaré group Clebsch-Gordan coefficients are the expansion coefficients of linear combination of tensor product states that transform irreducibly. The desired non-interacting irreducible states are computed by (1) constructing rest eigenstates of the two-body system, (2) decomposing them into irreducible

representations under $SU(2)$ rotations, (3) boosting the result to an arbitrary frame. The resulting Clebsch-Gordan coefficients in this basis are [33][19][32]

$$\begin{aligned} & \langle \mathbf{p}_1, \mu_1, \mathbf{p}_2, \mu_2 | (j, k) \mathbf{p}, \mu; l, s \rangle \\ &= \sum_{\mu_l \mu_s \mu'_1 \mu'_2} \delta(\mathbf{p} - \mathbf{p}_1 - \mathbf{p}_2) \frac{\delta(k - k(\mathbf{p}_1, \mathbf{p}_2))}{k^2} N^{-1}(p_1, p_2)(l, \mu_l, s, \mu_s | j, \mu)(j_1, \mu'_1, j_2, \mu'_2 | s, \mu_s) \times \\ & \quad Y_{l\mu_l}(\hat{\mathbf{k}}(\mathbf{p}_1, \mathbf{p}_2)) D_{\mu'_1 \mu_1}^{j_1} [R_w(B(\mathbf{p}/m_{120}), k_1)] D_{\mu'_2 \mu_2}^{j_2} [R_w(B(\mathbf{p}/m_{120}), k_2)] = \end{aligned} \quad (3)$$

$$\begin{aligned} &= \int d\hat{\mathbf{k}} \sum_{\mu_l \mu_s \mu'_1 \mu'_2} \delta(\mathbf{p}_1 - \mathbf{p}_1(\mathbf{p}, \mathbf{k})) \delta(\mathbf{p}_2 - \mathbf{p}_2(\mathbf{p}, \mathbf{k})) N(p_1, p_2)(l, \mu_l, s, \mu_s | j, \mu)(j_1, \mu'_1, j_2, \mu'_2 | s, \mu_s) \times \\ & \quad Y_{l\mu_l}(\hat{\mathbf{k}}) D_{\mu'_1 \mu_1}^{j_1} [R_w(B(\mathbf{p}/m_{120}), k_1)] D_{\mu'_2 \mu_2}^{j_2} [R_w(B(\mathbf{p}/m_{120}), k_2)]. \end{aligned} \quad (4)$$

In these expressions

$$p^\mu = p_1^\mu + p_2^\mu \quad m_{120}^2 = -p^2 \quad k^\mu = B^{-1}(\mathbf{p}/m_{120})^\mu{}_\nu \frac{1}{2}(p_1 - p_2)^\nu \quad (5)$$

$$N^{-2}(p_1, p_2) = \frac{\omega(k)\omega(k)(\omega(p_1) + \omega(p_2))}{\omega(p_1)\omega(p_2)(\omega(k) + \omega(k))} \quad (6)$$

and the two-body invariant mass

$$m_{120} := 2\sqrt{\mathbf{k}^2 + m^2} = 2\omega(k) \quad (7)$$

is replaced by the continuous variable $k := \sqrt{\mathbf{k}^2}$. The quantum numbers l and s are kinematically invariant quantities that distinguish multiple copies of representations with the same mass (k) and spin. For a two nucleon-system they have the same spectrum as the orbital and spin angular momentum operators in a partial wave representation of the non-relativistic basis.

An irreducible representation is also obtained by changing the order of the spin couplings in (3-4) where the orbital angular momentum is first coupled to one of the spins, $j_2 + l = I$, and then the result is coupled to the second spin. $j_1 + I = j$. This representation is constructed by making the replacements in

$$\begin{aligned} & \sum_{s, \mu_s} (j_1, \mu'_1, j_2, \mu'_2 | s, \mu_s)(l, \mu_l, s, \mu_s | j, \mu) \rightarrow \\ & \sum_{I, \mu_I} (l, \mu'_l, j_2, \mu_2 | I, \mu_I)(I, \mu_I, j_1, \mu'_1 | j, \mu) \end{aligned} \quad (8)$$

in equation (3) or (4). In this representation the degeneracy parameters (l, s) are replaced by (l, I) . When we construct three-body irreducible representations by successive pairwise coupling we use the coupling (3) in the first Clebsch-Gordan coefficient and the coupling (8) in the second Clebsch-Gordan coefficient. This allows us to identify the quantum numbers of the relativistic irreducible basis with the quantum numbers that we used in previous non-relativistic calculations ([34, 35]).

Three-particle irreducible representations for systems of non-interacting particles can be constructed by successive pairwise coupling of irreducible representations:

$$\begin{aligned} & \langle \mathbf{p}_1, \mu_1, \mathbf{p}_2, \mu_2, \mathbf{p}_3, \mu_3 | (J, q) \mathbf{P}, \mu; \lambda, I, j_{23}, k_{23}, l_{23}, s_{23} \rangle = \\ & \int d\mathbf{p}_{23} \sum_{\mu_{23}=-j_{23}}^{j_{23}} \langle \mathbf{p}_2, \mu_2, \mathbf{p}_3, \mu_3 | (j_{23}, k_{23}) \mathbf{p}_{23}, \mu_{23}; l_{23}, s_{23} \rangle \langle \mathbf{p}_1, \mu_1, \mathbf{p}_{23}, \mu_{23} | (J, q) \mathbf{P}, \mu; \lambda, I \rangle. \end{aligned} \quad (9)$$

For three nucleon scattering or bound state problems it is sufficient and convenient to work in the three-body center of momentum frame. This simplifies the coefficients; the Wigner rotations in the second Clebsch-Gordan coefficient in (9) become the identity and the normalization factor $N(p_{23}, p_1) \rightarrow 1$. Both of these factors are non-trivial in the first coefficient. The form of these coefficients for the case that particle 1 is a spectator (for details see [20]) and using a short hand notation, is

$$\begin{aligned} & \langle \mathbf{p}_1, \mu'_1, \mathbf{p}_2, \mu'_2, \mathbf{p}_3, \mu'_3 | (J, q) \mathbf{P} = \mathbf{0}, \mu; \lambda, I, j_{23}, k_{23}, l_{23}, s_{23} \rangle = \\ & \frac{\delta(\mathbf{0} - \mathbf{q}_1 - \mathbf{q}_2 - \mathbf{q}_3) \frac{1}{N(q_2, q_3)}}{\frac{\delta(q_1 - q) \delta(k(\mathbf{q}_2, \mathbf{q}_3) - k)}{q^2 k^2}} \\ & \sum_{\mu_2 \mu_3 \mu_s} \sum_{\mu_1 \mu_\lambda \mu_I} \left(\frac{1}{2}, \mu_2, \frac{1}{2}, \mu_3 | s, \mu_s \right) (l, \mu_l, s, \mu_s, | j, \mu_j) (\lambda, \mu_\lambda, \frac{1}{2}, \mu'_1 | I, \mu_I) (j, \mu_j, I, \mu_I | J, \mu) \\ & Y_{\mu_\lambda}^\lambda(\hat{\mathbf{q}}_1) Y_{\mu_l}^l(\hat{\mathbf{k}}(\mathbf{q}_2, \mathbf{q}_3)) \\ & D_{\mu'_2 \mu_2}^{\frac{1}{2}} [R_w(B(-\mathbf{q}_1/m_{023}), k_2(\mathbf{q}_2, \mathbf{q}_3))] D_{\mu'_3 \mu_3}^{\frac{1}{2}} [R_w(B(-\mathbf{q}_1/m_{023}), k_3(\mathbf{q}_2, \mathbf{q}_3))]. \end{aligned} \quad (10)$$

where

$$(\mathbf{q}_i, \omega(q_i)) = B^{-1}(\mathbf{P}/M) p_i, \quad i \in \{1, 2, 3\} \quad (\mathbf{k}_i, \omega(k_i)) = B^{-1}(-\mathbf{q}_k/m_{ij}) p_i \quad (11)$$

$$\sum_i \mathbf{q}_i = \mathbf{0} \quad M = \sum_{i=1}^3 \sqrt{m^2 + \mathbf{q}_i^2} \quad (12)$$

The important property of the states

$$|(j, k) \mathbf{p}, \mu; l, s\rangle \quad (13)$$

and

$$|(J, q)\mathbf{P}, \mu; \lambda, I, j_{23}, k_{23}, l_{23}, s_{23}\rangle \quad (14)$$

is that they transform irreducibly. The mass and spin are given by

$$m_{120} = 2\sqrt{m^2 + \mathbf{k}^2} \quad j \quad (15)$$

$$M = \sqrt{4m^2 + 4\mathbf{k}^2 + \mathbf{q}^2} + \sqrt{m^2 + \mathbf{q}^2} \quad J \quad (16)$$

respectively.

In order to use these representations to construct dynamical representations an interaction is added to the two- or three-body invariant mass operator of the form

$$\langle (j, k)\mathbf{p}, \mu \cdots |v| \cdots \mathbf{p}', \mu'(j', k') \rangle = \delta_{\mu\mu'} \delta_{jj'} \delta(\mathbf{p} - \mathbf{p}') \langle k, \cdots \|v^j\| \cdots', k' \rangle \quad (17)$$

for $N = 2$ or

$$\langle (J, q)\mathbf{P}, \mu \cdots |V| \cdots \mathbf{P}', \mu'(J', q') \rangle = \delta_{\mu\mu'} \delta_{JJ'} \delta(\mathbf{P} - \mathbf{P}') \langle k, q, \cdots \|V^J\| \cdots', k', q' \rangle \quad (18)$$

for $N = 3$. Diagonalizing $m_{12} = m_{120} + v$ or $M = M_0 + V$ in the noninteracting irreducible basis gives simultaneous eigenstates of $m_{12}, \mathbf{p}, j^2, j_z$ for $N = 2$ and of M, \mathbf{P}, J^2, J_z for $N = 3$. In both the two and three-body case these eigenstates, $|(j_{12}, \lambda_{m_{12}})\mathbf{p}_{12}, \mu_{12} \cdots \rangle$ and $|(J, \lambda_M)\mathbf{P}, \mu \cdots \rangle$, where $\lambda_{m_{12}}$ and λ_M are the eigenvalues of m_{12} and M , are complete on the two and three-body Hilbert spaces respectively.

The dynamical representation of the Poincaré group is defined by requiring that these eigenstates transform like (1) with the mass being replaced by the mass eigenvalues λ_M or $\lambda_{m_{12}}$. This representation is unitary and defines the dynamics of the system. With our choice of irreducible basis, $\{\mathbf{p}, j_{cz}\}$, the resulting irreducible representations of the Poincaré group have a mass independent representation of the three-dimensional Euclidean subgroup, which Dirac [36] called an “instant-form dynamics”.

For the three-nucleon case there remains the problem of how to construct realistic NN -interactions. For two-body interactions the relation

$$H_{12}^2 - \mathbf{p}^2 = m_{12}^2 = 4(\mathbf{k}^2 + m^2) + 4mv_{NN} = 4m \underbrace{(\mathbf{k}^2/m + v_{NN} + m)}_{h_{nr} = H_{nr} - \frac{\mathbf{p}^2}{4m}} \quad (19)$$

implies that the square of the two-body mass operator has a simple relation to the non-relativistic rest Hamiltonian with a “realistic” NN interaction [25] provided one identifies the spectrally equivalent relative momenta, \mathbf{k}^2 . In the relativistic case $k := B^{-1}(\mathbf{p}/m_{120})^{\frac{1}{2}}(p_1 - p_2)$

$$\begin{aligned}\mathbf{k} &\equiv \mathbf{k}(\mathbf{p}_1, \mathbf{p}_2) \\ &= \frac{1}{2}(\mathbf{p}_1 - \mathbf{p}_2 - \mathbf{P} \frac{\omega(p_2) - \omega(p_1)}{\omega(p_2) - \omega(p_1) + \sqrt{(\omega(p_2) - \omega(p_1))^2 - \mathbf{P}^2}}),\end{aligned}\quad (20)$$

while in the non relativistic case $k = B_g^{-1}(\mathbf{p}/2m)^{\frac{1}{2}}(p_1 - p_2)$:

$$\mathbf{k} = \frac{1}{2} \left(\left(\mathbf{p}_1 - \frac{\mathbf{P}}{2m} m \right) - \left(\mathbf{p}_2 - \frac{\mathbf{P}}{2m} m \right) \right) = \frac{1}{2}(\mathbf{p}_1 - \mathbf{p}_2) \quad (21)$$

where $B(\mathbf{p}/m_{120})$ is a rotationless Lorentz boost and $B_g(\mathbf{p}/2m)$ is the corresponding Galilean boost.

The Kato-Birman invariance principle [37–39] implies that the Møller wave operators satisfy

$$\Omega_{\pm}(H, H_0) := s - \lim_{t \rightarrow \pm\infty} e^{iHt} e^{-iH_0 t} = s - \lim_{t \rightarrow \pm\infty} e^{if(H)t} e^{-if(H_0)t} = \Omega_{\pm}(f(H), f(H_0)) \quad (22)$$

where $f(x)$ is any piecewise differentiable function of bounded variation with positive derivative [39]. The functions $f(x) = x^2$ and $f(x) = x^{1/2}$ satisfy the conditions of Kato-Birman theorem. Using equation (22) along with the kinematic Euclidean invariance of the Hamiltonians H_r and H_{nr} gives the following relation between the two-body scattering wave operators

$$\Omega_{\pm}(H_r, H_{r0}) = \Omega_{\pm}(H_r^2, H_{r0}^2) = \Omega_{\pm}(M_r^2, M_{r0}^2) = \Omega_{\pm}(M_r, M_{r0}). \quad (23)$$

On the other hand the identification (19) along with the reparametrization $t \rightarrow t' = 4mt$ gives

$$\Omega_{\pm}(M_r^2, M_{r0}^2) = s - \lim_{t \rightarrow \pm\infty} e^{ihn_r 4mt} e^{-h_{nr0} 4mt} = s - \lim_{t' \rightarrow \pm\infty} e^{ihn_r t'} e^{-h_{nr0} t'} = \Omega_{\pm nr}(h_{nr}, h_{nr0}). \quad (24)$$

Writing both wave operators as direct integrals over $\mathbf{k}^2 = mh_{0nr} = (M_0^2 - 4m^2)/4$ leads to the identifications

$$\Omega_{\pm}(H_{nr}, H_{0nr}) = \Omega_{\pm}(h_{nr}, h_{0nr}) = \int_{\oplus} \hat{\Omega}_{\pm}(\mathbf{k}^2) d\mathbf{k}^2 = \Omega_{\pm}(H_r, H_{0r}) = \Omega_{\pm}(M_{nr}^2, M_{0nr}^2) \quad (25)$$

and

$$S(\mathbf{k}^2) = \Omega_{r+}^{\dagger}(\mathbf{k}^2) \Omega_{r-}(\mathbf{k}^2) = \Omega_{nr+}^{\dagger}(\mathbf{k}^2) \Omega_{nr-}(\mathbf{k}^2). \quad (26)$$

The identification of the relativistic and non-relativistic wave operators as functions of \mathbf{k}^2 ensures that the relativistic two-body model is fit to the same two-body S -matrix (experimental data) as the non-relativistic model provided the interactions are related by (19). The identification of the

wave operators also implies the identity of the scattering wave functions as a function of \mathbf{k} . The identity of the bound state wave functions is also due to the relation (19).

In our calculations we use the interaction v defined by $v := m_{12} - m_{120}$ which we construct [40] from the NN interaction in (19) by iterating

$$\{m_{120}, v\} = 4mv_{NN} - v^2 \quad (27)$$

in the irreducible plane-wave basis. Because m_{12} and m_{12}^2 have the same eigenvectors, the \mathbf{k} dependence of the wave functions constructed from m_{12} are also identical to the corresponding non-relativistic wave functions.

The equation (27) in the irreducible plane-wave basis has the form

$$\begin{aligned} \langle k, l, s | v^j | k', l', s' \rangle = \\ 2m \frac{\langle k, l, s | v_{NN}^j | k', l', s' \rangle}{\omega(k) + \omega(k')} - \sum_{l'' s''} \int k''^2 dk'' \frac{\langle k, l, s | v^j | k'', l'', s'' \rangle \langle k'', l'', s'' | v^j | k', l', s' \rangle}{2\omega(k) + 2\omega(k')} \end{aligned} \quad (28)$$

The iteration converges quickly for realistic interactions [40]. While a mathematical proof of convergence of the iteration of (28) is lacking, the results of the iterations are easily tested because the resulting $m_{120} + v$ must have the same eigenfunctions as the non-relativistic two-body Hamiltonian.

We applied this approach using the CD Bonn potential as the non-relativistic interaction $v_{NN}(k, k')$. In our previous studies [20, 27, 28] we used the momentum transformation of ref. [24] and in addition restricted to leading order terms in \mathbf{p}/ω and v/ω expansion only

$$V(\mathbf{k}, \mathbf{k}'; \mathbf{q}) = v(\mathbf{k}, \mathbf{k}') \left(1 - \frac{\mathbf{q}^2}{8\omega(k)\omega(k')} \right). \quad (29)$$

We checked that in most cases this simple approximation leads to practically the same results as the exact approach applied in the present study.

Once the two-body mass operator is constructed, the three-body mass operator for the interacting (ij) pair is the well-defined non-linear function of the two-body mass:

$$M_{(ij)(k)} = \sqrt{(m_{ij0} + v_{ij})^2 + \mathbf{q}_k^2} + \sqrt{m^2 + \mathbf{q}_k^2}. \quad (30)$$

where v_{ij} is embedded in the three body Hilbert space so it commutes with \mathbf{q}_k . If this is interpreted as the rest energy operator, the interacting pair and spectator energies are additive in the rest frame. This implies that the S matrix clusters properly in the rest frame - while the invariance of S in all frames ensures that this property extends to all inertial coordinate systems.

Pairwise interactions in the three-body system are defined by

$$V_{(ij)(k)} = M_{(ij)(k)} - M_0 = \sqrt{(m_{ij0} + v_{ij})^2 + \mathbf{q}^2} - \sqrt{m_{ij0}^2 + \mathbf{q}^2}. \quad (31)$$

For any pair of particles these interactions commute with kinematic momentum and spin, and are independent on the momentum and magnetic quantum numbers. This ensures that the sum of the interactions has the general form (18).

A generalization of the method used in (27-28) can be used to construct $V_{(ij)(k)}$ by iterating

$$\{\sqrt{m_{ij0}^2 + \mathbf{q}^2}, V_{(ij)(k)}\} = v^2 + \{m_{ij0}, v\} - V_{(ij)(k)}^2. \quad (32)$$

Specifically

$$\begin{aligned} \langle k, \dots | V_{(ij)(k)}(\mathbf{q}^2) | k', \dots \rangle &= \frac{1}{\sqrt{m_{ij0}^2(k) + \mathbf{q}^2} + \sqrt{m_{ij0}^2(k') + \mathbf{q}^2}} \times \\ & \left[\langle k, \dots | v^2 | k', \dots \rangle + 2(\omega(k) + \omega(k')) \langle k | v | k' \rangle \right. \\ & \left. - \sum_{l''s''} \int \langle k, \dots | V_{(ij)(k)}(\mathbf{q}^2) | k'', \dots \rangle k''^2 dk'' \langle k'', \dots | V_{(ij)(k)}(\mathbf{q}^2) | k', \dots \rangle \right] \end{aligned} \quad (33)$$

This iteration also converges and is used to construct interactions V_{ij} for each pair of particles.

The three-body mass (rest energy) operator is

$$M = M_0 + V_{12} + V_{23} + V_{31} \quad (34)$$

where M_0 is the three-body kinematic invariant mass:

$$M_0 = \sqrt{m_{120}^2 + \mathbf{q}^2} + \sqrt{m^2 + \mathbf{q}^2} \quad (35)$$

Our relativistic Faddeev equation is based on this mass operator (34) with two-body interactions constructed from the CD Bonn interaction using equations (27), (28), (32) and (33).

Finally, just like in the non-relativistic case, there is a natural order of coupling of the irreducible representations for computing each pairwise interaction. The change of basis relating different orders of coupling is needed for the implementation of the Faddeev equation as an integral equation. The required basis change only changes the invariant degeneracy quantum numbers associated with each order of coupling.

$$\langle (jm)\mathbf{P}, \mu(ab)(c) | (j'm')\mathbf{P}', \mu'(de)(f) \rangle = \delta(\mathbf{P} - \mathbf{P}') \delta_{jj'} \delta_{\mu\mu'} R^{jm}[(ab)(c); (de)(f)] \quad (36)$$

The invariants $R^{jm}[(ab)(c); (de)(f)]$ are Racah coefficients for the Poincaré group. They are constructed using four Poincaré Clebsch-Gordan coefficients. We compute this quantity using the Balian-Brezin method [41], where the variables associated with one order of coupling are expressed in terms of the variables associated with another order of coupling. The invariant coefficient, $R^{jm}[(ab)(c); (de)(f)]$, can be computed by evaluating the expression at zero momentum, averaging over the magnetic quantum numbers, and evaluating the resulting expression at any kinematically allowed set of momenta [42, 43]. These Racah coefficients contain Wigner rotations and jacobians which do not appear in the non-relativistic permutation operators. Explicit expressions are given in Appendix B.

III. SPIN OBSERVABLES

In a Poincaré invariant quantum theory or relativistic quantum field theory the spin of a particle can be defined as the angular momentum that is measured in the particle's rest frame. For any non-zero momentum, \mathbf{p} , different Lorentz transformations can be used to transform a particle at rest to frame where the particle has momentum \mathbf{p} . Because the commutator of two rotationless boost generators

$$[K_j, K_k] = -i\epsilon_{jkl}J_l \quad (37)$$

is a rotation generator, the spin of the particle with momentum \mathbf{p} depends on the choice of Lorentz transformation that transforms the particle's momentum from zero to \mathbf{p} . In order to have an unambiguous definition of the spin it is necessary to choose a standard set of \mathbf{p} dependent Lorentz boosts, $B(\mathbf{p}/m)^\mu{}_\nu$, that transform a particle of mass m at rest to momentum \mathbf{p} . Then the spin of the particle can be unambiguously defined as the value of the spin measured in the particles rest frame if it is transformed to the rest frame using the standard Lorentz transformation. With this definition, if two spins are equal in one frame they are equal in all frames.

The choice of standard boost is not unique because if $R(\mathbf{p}/m)$ is any \mathbf{p} dependent rotation and

$$B'(\mathbf{p}/m) := B(\mathbf{p}/m)R(\mathbf{p}/m) \quad (38)$$

then both $B^{-1}(\mathbf{p}/m)^\mu{}_\nu$ and $B'^{-1}(\mathbf{p}/m)^\mu{}_\nu$ both transform \mathbf{p} to zero.

Each choice of boost leads to a different spin operator, corresponding to a different prescription for measuring the spin in an arbitrary frame. The rotation

$$R(\mathbf{p}/m) = B^{-1}(\mathbf{p}/m)B'(\mathbf{p}/m) \quad (39)$$

that relates different boosts is called a generalized Melosh rotation [44][32].

The spin operator \mathbf{j}_x associated with a boost $B_x(\mathbf{p}/m)$ is defined as the following function of the Poincaré generators:

$$(0, \mathbf{j}_x) = B_x^{-1}(\mathbf{p}/m)^\mu{}_\nu W^\nu / m \quad (40)$$

where $W^\mu = \frac{1}{2}\epsilon^{\mu\alpha\beta\gamma}p_\alpha M_{\beta\gamma}$ is the Pauli-Lubanski vector, $M_{\beta\gamma}$ is the relativistic angular momentum tensor, and $B_x(\mathbf{p}/m)$ is the boost matrix with the parameters \mathbf{p}/m and m replaced by the mass and momentum operators.

While this quantity has the appearance of a four vector, it is not because of the operator dependence of the arguments of the boost. Instead, under Lorentz transformations the spin transforms like

$$U^\dagger(\Lambda, 0)\mathbf{j}_x U(\Lambda, 0) = R_{wx}(\Lambda, p)\mathbf{j}_x \quad (41)$$

where

$$R_{wx}(\Lambda, p) := B_x^{-1}(\Lambda p)\Lambda B_x(\mathbf{p}) \quad (42)$$

is the Wigner rotation associated with the boost $B_x(\mathbf{p}/m)$. It is a consequence of the Poincaré commutation relations that the components of any of these spin observables satisfy $SU(2)$ commutation relations

$$[j_{xl}, j_{xm}]_- = i\epsilon_{lmn}j_{xn} \quad (43)$$

for any “ x ”. The operator \mathbf{j}^2 is independent of the choice of boost because the generalized Melosh rotations leave the scalar product of two vectors unchanged. The spin defined with the textbook rotationless or canonical boost is called the canonical spin.

It is natural to ask how these different types of spins are measured in the laboratory. Spins of isolated elementary or composite particles are measured in the laboratory through their response to classical electromagnetic fields. In the one-photon exchange approximation the photon couples to matrix elements of a covariant current operator. Imposing Poincaré covariance, current conservation, and discrete symmetries allows one to express all current matrix elements in terms

of an independent set of matrix elements, which have a 1-1 correspondence with invariant form factors. All conventional form factors can be expressed in terms of Breit frame matrix elements with canonical spin and a quantization axis parallel to the Breit frame momentum transfer.

In the $SL(2, C)$ representation canonical boosts are represented by positive Hermitian matrices. They have the general form

$$B(\mathbf{p}/m) = \exp(\boldsymbol{\sigma} \cdot \boldsymbol{\rho}/2) \quad (44)$$

where $\boldsymbol{\rho}$ is the rapidity of the Lorentz transformation and $\boldsymbol{\sigma}$ are the Pauli matrices. In this paper all of our spins are canonical spins. The $SO(1, 3)$ representation of canonical boosts are the standard rotationless boosts.

Given a definition of the form factors in terms of independent current matrix elements in a given basis, it is also possible to express them in terms of current matrix elements in any other standard frame using any other basis [45]. For example the expression in terms of Breit frame canonical spin matrix elements, can be replaced by a different independent set of laboratory frame helicity spin matrix elements. In quantum field theory the choice of boost is built into conventions used to define the Dirac spinors. The relation of the invariant form factors to current matrix elements with different choices of spin determines the relationship between different spin observables and experiment.

The spin degrees of freedom of the asymptotic incoming or outgoing particles are most conveniently expressed in terms of traces of density matrices, which is a reflection of the fact that realistic initial and/or final states are generally not pure states. Scattering spin observables in cross sections are formally defined by [34]

$$\langle O \rangle = \frac{\text{Tr}(S_f T S_i T^\dagger)}{\text{Tr}(T T^\dagger)} \quad (45)$$

where T is the invariant scattering amplitude for the reaction under consideration. The connection between the invariant scattering amplitude defined in the particle data book [46] and the transition amplitudes constructed by solving our formulation of the relativistic Faddeev equations is given in [47].

The quantities S have the form

$$S = \sum s_a S_a \quad (46)$$

where the index runs over all N_i or N_f initial or final sets of magnetic quantum numbers, S_a are a basis for $N_i \times N_i$ or $N_f \times N_f$ matrices that are orthonormal with respect to the trace norm and s_a are constant coefficients [34].

If the initial and final asymptotic states are represented in a canonical spin basis, then the magnetic quantum numbers that appear in the invariant amplitudes T transform with Wigner rotations under Lorentz transformations. The result is that the spin observable $\langle O \rangle$ will not be invariant unless the matrices S_a or coefficients s_a are defined to transform in a manner that leaves the observable invariant.

Any spin observable can be made Lorentz invariant, by defining the invariant observable as its value in a given frame if it is transformed to the frame with a specific Lorentz boost. This can be used to get an invariant definition of the vector or tensor polarizations.

In this paper invariant spin observables are defined to be the values of the observable in the laboratory frame (rest frame of the target). To evaluate the corresponding spin observable it is only necessary to evaluate the expression

$$\langle O \rangle = \frac{\text{Tr}(S_f T S_i T^\dagger)}{\text{Tr}(T T^\dagger)} \quad (47)$$

for the values of the invariant amplitudes with laboratory kinematics.

This observable is equal its value evaluated in other frames using the formula

$$\langle O \rangle = \frac{\text{Tr}(S_f D^\dagger M D' S_i D'^\dagger M^\dagger D)}{\text{Tr}(M M^\dagger)} \quad (48)$$

where the invariant amplitudes are evaluated in the other frame and D and D' are products of Wigner D -functions of the Wigner rotations associated with the boost from the lab frame to the other frame.

Our specific interest in this paper is the observable A_y with polarized incoming nucleon. The convention used to define A_y is the Madison convention, where the laboratory frame scattering plane is in the xz plane. The observable A_y , is defined as

$$\langle A_y \rangle = \frac{\text{Tr}(T(\sigma_y \times I_d)T^\dagger)}{\text{Tr}(T T^\dagger)} \quad (49)$$

Because the Wigner rotation for canonical boost along the direction of a particle's momentum is the identity, σ_y is unchanged so A_y can also be evaluated in the c.m. frame without making any compensating Wigner rotations.

IV. FADDEEV EQUATION

The nucleon-deuteron scattering with neutron and protons interacting through a NN interaction v_{NN} alone is described in terms of a breakup operator T satisfying the Faddeev-type integral equation [34, 48]

$$T|\phi\rangle = tP|\phi\rangle + tPG_0T|\phi\rangle. \quad (50)$$

The two-nucleon (2N) t -matrix t results from solving the Lippmann-Schwinger equation with the interaction v_{NN} . The permutation operator $P = P_{12}P_{23} + P_{13}P_{23}$ is given in terms of the transposition P_{ij} which interchanges nucleons i and j . The incoming state $|\phi\rangle = |\mathbf{q}_0\rangle|\phi_d\rangle$ describes the free nucleon-deuteron motion with relative momentum \mathbf{q}_0 and the deuteron state vector $|\phi_d\rangle$. Finally G_0 is resolvent of the three-body center of mass kinetic energy.

The elastic nd scattering transition operator U is given in terms of T by [34, 48]

$$U = PG_0^{-1} + PT. \quad (51)$$

This is our standard non-relativistic formulation, which is equivalent to the non-relativistic 3N Schrödinger equation plus boundary conditions. The formal structure of these equations in the relativistic case remains the same but the ingredients change. As explained in [22] the relativistic 3N rest Hamiltonian (mass operator) has the same form as the non-relativistic one, only the momentum dependence of the kinetic energy changes and the relation of the pair interactions in the three-body problem to the pair interactions in the two-body problem changes. Consequently all the formal steps leading to Eqs.(50) and (51) remain the same.

The free relativistic invariant mass of three identical nucleons in their c.m. system has the form (35) while the free two-body mass operator has the form (7).

As introduced in [19] the pair forces in the relativistic 3N $2 + 1$ mass operator are given by (31) where $V = V(\mathbf{q}^2)$ reduces to the interaction v for $\mathbf{q} = 0$.

The transition matrix that appears in the kernel of the Faddeev equation (50) is obtained by solving the ‘‘Lippmann-Schwinger equation’’, which must be solved as a function of \mathbf{q}^2

$$t(\mathbf{k}, \mathbf{k}'; \mathbf{q}^2) = V(\mathbf{k}, \mathbf{k}'; \mathbf{q}^2) + \int d^3k'' \frac{V(\mathbf{k}, \mathbf{k}''; \mathbf{q}^2)t(\mathbf{k}'', \mathbf{k}'; \mathbf{q}^2)}{\sqrt{(2\omega(\mathbf{k}')^2 + \mathbf{q}^2) - \sqrt{(2\omega(\mathbf{k}'')^2 + \mathbf{q}^2) + i\epsilon}}. \quad (52)$$

The input two-body interactions are computed by solving equation (32) and (33).

The new relativistic ingredients in Eqs.(50) and (51) will therefore be the t -operator (52) (expressed in partial waves) and the resolvent of the 3N invariant mass

$$G_0 = \frac{1}{E + i\epsilon - M_0}, \quad (53)$$

where M_0 is given by Eq. (35) and E is the total 3N c.m. energy expressed in terms of the initial neutron momentum \mathbf{q}_0 relative to the deuteron

$$E = \sqrt{(M_d)^2 + \mathbf{q}_0^2} + \sqrt{m^2 + \mathbf{q}_0^2}, \quad (54)$$

and M_d is the deuteron rest mass.

Currently Eq.(50) in its non-relativistic form is numerically solved for any NN interaction using a momentum space partial-wave decomposition. Details are presented in ref. [48]. This turns Eq.(50) into a coupled set of two-dimensional integral equations. As shown in [20], in the relativistic case we can keep the same formal structure, though the permutation operators are replaced by the corresponding Racah coefficients (36) for the Poincaré group. These coefficients include both jacobians and Wigner rotations that do not appear in the non-relativistic permutation operators [34, 35]. These coefficients are computed in the appendix, using methods that we have applied to compute the non-relativistic permutation operators.

In the non-relativistic case the partial wave projected momentum space basis is

$$|pq(ls)j(\lambda\frac{1}{2})IJ(t\frac{1}{2})T\rangle, \quad (55)$$

where p and q are the magnitudes of standard Jacobi momenta (see [34, 35]), obtained by transforming single particle momenta to the rest frame of a two or three-body system using Galilean boosts, and $(ls)j$ two-body quantum numbers with obvious meaning, $(\lambda 1/2)I$ refer to the third nucleon (described by the momentum q), J is the total 3N angular momentum and the rest are isospin quantum numbers. In the relativistic case this basis is replaced by the irreducible plane wave states defined in (10).

The basis states (10) are used for the evaluation of the partial wave representation of the permutation operator P with Wigner rotations of spin states for nucleons 2 and 3 included. In the relativistic case we adopt the following short-hand notation for the irreducible three-body states, which also includes isospin quantum numbers coupled in the same order:

$$|k, q, \alpha\rangle := |kq(ls)j(\lambda, \frac{1}{2})IJ(t\frac{1}{2})T\rangle = |(J, q)\mathbf{P}, \mu; \lambda, I, j_{23}, k_{23}, l_{23}, s_{23})|(t\frac{1}{2})T\rangle \quad (56)$$

Equipped with that, projecting Eq.(50) onto the basis states $|k, q, \alpha\rangle$ one encounters like in the non-relativistic notation [35]

$${}_1\langle kq\alpha|P|k'q'\alpha'\rangle_1 = {}_1\langle kq\alpha|k'q'\alpha'\rangle_2 + {}_1\langle kq\alpha|k'q'\alpha'\rangle_3 = 2 {}_1\langle kq\alpha|k'q'\alpha'\rangle_2. \quad (57)$$

This is evaluated by inserting the complete basis of states $|\mathbf{p}_1, \mu_1, \mathbf{p}_2, \mu_2, \mathbf{p}_3, \mu_3\rangle$ and using (10). It can be expressed in a form which resembles closely the one appearing in the non-relativistic regime [34, 35]

$${}_1\langle k q \alpha | P | k' q' \alpha' \rangle_1 = \int_{-1}^1 dx \frac{\delta(k - \pi_1)}{k^2} \frac{\delta(k' - \pi_2)}{k'^2}$$

$$\frac{1}{N_1(q, q', x)} \frac{1}{N_2(q, q', x)} G_{\alpha\alpha'}^{BB}(q, q', x), \quad (58)$$

where all ingredients are given in the Appendix.

Due to short-range nature of the NN interaction it can be considered negligible beyond a certain value j_{max} of the total angular momentum in the two-nucleon subsystem. Generally with increasing energy j_{max} will also increase. For $j > j_{max}$ we put the t-matrix to be zero, which yields a finite number of coupled channels for each total angular momentum J and total parity $\pi = (-)^{l+\lambda}$ of the 3N system. To achieve converged results at our energies we used all partial wave states with total angular momenta of the 2N subsystem up to $j_{max} = 5$ and took into account all total angular momenta of the 3N system up to $J = 25/2$. This leads to a system of up to 143 coupled integral equations in two continuous variables for a given J and parity.

V. RESULTS AND DISCUSSION

The subject of the present study is to investigate the influence of relativity on the nd elastic scattering nucleon analyzing power A_y at low energies. We define the invariant observable A_y to be the value measured in the laboratory frame (target at rest).

To this aim we solved Faddeev equations at a number of the incoming neutron lab energies $E_n = 5$ MeV, 8.5 MeV, and 13 MeV. To check the energy dependence of the effect we added two additional energies $E_n = 35$ MeV and 65 MeV. In order to see the importance of specific relativistic features we solved the equation in the relativistic case with and without Wigner rotations. This allowed us to see which effects, dynamical corrections, which are induced by the momentum dependence of the two-body force together with kinematical relativistic effects coming from the use of the Poincare Jacobi variables, or Wigner rotations, dominate for A_y .

Figs.1 and 2 illustrate the results. When only dynamical effects are taken into account (Wigner rotations neglected) then at low energies of the incoming neutron the relativistic and non relativistic predictions are practically the same with the exception of the angular region close to the maximum of A_y where the relativistic prediction is $\approx 2\%$ below the non-relativistic one (see Fig.1). This small effect disappears at higher energies (see Fig.2). Including Wigner rotations lowers significantly the values of A_y in a large region of angles around the maximum. The changes in the maximum are up to $\approx 10\%$. Again, when the energy of the incoming neutron increases non relativistic and relativistic predictions are practically identical.

The large changes of A_y occur in a region of energies where this observable is extremely sensitive to changes in 3P_0 , 3P_1 , and ${}^3P_2 - {}^3F_2$ NN force components [49]. At energies where this sensitivity

dies out also the relativistic effects for A_y become negligible. This allows to conclude that the large effects seen for A_y at low energies are due to amplification of changes of the 3P_j contributions due to relativity by a large sensitivity of A_y to P -waves.

In a recent study [29] the changes of A_y due to relativity by 10% at $E_n^{lab} = 3$ MeV have been reported. Very probably the opposite sign of the effect found in that study can be attributed to the impulse approximation used when calculating A_y .

VI. SUMMARY AND OUTLOOK

We numerically solved the 3N Faddeev equation for nd scattering including relativistic kinematics, dynamical relativistic effects and Wigner rotations at the neutron lab energies $E_n^{lab} = 5, 8.5, 13, 35,$ and 65 MeV. As dynamical input we took the non-relativistic NN potential CD Bonn and generated in the 2N c.m. system an exactly on-shell equivalent relativistic interaction v , by solving numerically nonlinear quadratic equation relating matrix elements of the non-relativistic and relativistic potentials. We checked that the approximate procedure using an analytical scale transformation of momenta applied in our previous studies provides practically the same results as the present exact approach. In addition a similar nonlinear equation (Eq.(32)) was used to generate the momentum dependent two-body interaction embedded in the 3-particle Hilbert space.

We found that at low energies the effects of Wigner rotations are most important for the analyzing power. They lower the maximum of A_y by up to $\approx 10\%$. The dynamical relativistic effects are of minor importance for A_y and provide small changes of A_y in a region close to its maximum. They lower A_y by only $\approx 2\%$. The relativistic effects disappear at higher energies.

Wigner rotations are negligible for the cross section and all other spin observables in elastic Nd scattering with exception of four low energy spin correlations $C_{z,yz}, C_{x,xy}, C_{y,xx-yy}, C_{y,yy}$ and four low energy spin transfer coefficients: from deuteron to deuteron $K_z^{y'z'}$ and $K_x^{x'y'}$, and from deuteron to neutron $K_{yz}^{z'}$ and $K_{xy}^{x'}$. Very probably this can be traced back as in the case of A_y to their sensitivity to 3P -waves.

These results shed new light on the low-energy analyzing power puzzle. It is known that the existing discrepancies between A_y data and theoretical predictions based on NN potentials only cannot be removed when current three-nucleon force, mostly of 2π -exchange character [10, 11] are included in the nuclear Hamiltonian. This indicated that additional 3N forces should be added to the 2π -exchange type forces. Such forces provided by χ PT in NNLO and NNNLO orders are expected to provide the solution for the A_y puzzle [9, 54]. It seems that in view of the present




FIG. 1: (color online) The nucleon analyzing power A_y for nd elastic scattering at various lab energies E_n^{lab} of the incoming neutron. The dotted line is the result of the non-relativistic Faddeev calculation with the CD Bonn potential. The relativistic predictions without and with Wigner spin rotations are shown by the dashed and solid lines, respectively. The nd data at 5 MeV and 8.5 MeV are from ref. [50] and at 13 MeV are from ref. [51].




FIG. 2: (color online) The nucleon analyzing power A_y for nd elastic scattering at $E_n^{lab} = 35$ MeV and 65 MeV. For description of lines see Fig.1. All theoretical predictions are practically overlapping. The pd data at 35 MeV are from ref. [52] and at 65 MeV from [53].

result they must increase the maximum of A_y stronger than expected up to now. However, A_y is a very sensitive observable and our approach, using the irreducible $\{\mathbf{p}, j_{cz}\}$ basis, is only one of many possible basis choices that lead to Poincaré invariant dynamical theories that are two-body

scattering equivalent, which may give different three-body predictions.

Acknowledgments

The numerical calculations were performed on the IBM Regatta p690+ of the NIC in Jülich, Germany. This work was performed in part under the auspices of the U. S. Department of Energy, Office of Nuclear Physics, contract No. DE-FG02-86ER40286 with the University of Iowa.

APPENDIX A: SPINORS

In our calculations $SL(2, \mathbb{C})$ matrices are used to represent Wigner rotations and Lorentz transformations. In this way spin algebra is reduced to working with 2×2 complex matrices, and there is no need for representations in terms of Euler angles. The relevant relations are give below.

The coordinates of a four-vector p^μ can be labeled by the 2×2 Hermitian matrix P :

$$P := p^\mu \sigma_\mu = \begin{pmatrix} p^0 + p^3 & p^1 - ip^2 \\ p^1 + ip^2 & p^0 - p^3 \end{pmatrix}, \quad (\text{A1})$$

where σ_μ are the identity and the three Pauli matrices. The components of p^μ can be extracted from the matrix P using:

$$p^\mu = \frac{1}{2} \text{Tr}(\sigma_\mu P) = \frac{1}{2} \text{Tr}(P \sigma_\mu). \quad (\text{A2})$$

Because

$$\det(P) = (p^0)^2 - (\vec{p})^2 = -\eta_{\mu\nu} p^\mu p^\nu = m^2 \quad (\text{A3})$$

and

$$P = P^\dagger \quad (\text{A4})$$

for real p^μ , it follows that any linear transformation that preserves the Hermiticity and determinant of P is a real Lorentz transformation. It is easy to show that if A is a complex 2×2 matrix with $\det(A) = 1$ then the transformation

$$P \rightarrow P' = APA^\dagger \quad (\text{A5})$$

has both of these properties:

$$\det(P') = \det(P) \quad \text{and} \quad P = P^\dagger \rightarrow P' = P'^\dagger. \quad (\text{A6})$$

The most general 2×2 matrix with determinant 1 can be written as

$$A = \pm e^{\frac{i}{2} \boldsymbol{\sigma} \cdot \mathbf{z}} \quad (\text{A7})$$

where $\mathbf{z} = \boldsymbol{\theta} - i\rho$. If $\rho = \mathbf{0}$ then $A = U(\boldsymbol{\theta})$ is unitary and corresponds to an $SU(2)$ rotation through an angle $|\boldsymbol{\theta}|$ about the $\hat{\boldsymbol{\theta}}$ axis. If $\boldsymbol{\theta} = \mathbf{0}$ then A is a positive Hermitian matrix that corresponds to a rotationless (canonical) Lorentz boost with rapidity $|\rho|$ in the direction $\hat{\boldsymbol{\rho}}$.

The rotation $U(\boldsymbol{\theta})$ is given by

$$A \rightarrow U(\boldsymbol{\theta}) = \sigma_0 \cos\left(\frac{\theta}{2}\right) + i\boldsymbol{\sigma} \cdot \hat{\boldsymbol{\theta}} \sin\left(\frac{\theta}{2}\right). \quad (\text{A8})$$

The axis of rotation can be extracted using

$$\hat{\boldsymbol{\theta}} = -i \frac{\text{Tr}(\boldsymbol{\sigma} U(\boldsymbol{\theta}))}{|\text{Tr}(\boldsymbol{\sigma} U(\boldsymbol{\theta}))|} \quad (\text{A9})$$

and the angle of rotation can be extracted from

$$\theta = 2 \tan^{-1} \left(\frac{\text{Tr}(-i\boldsymbol{\sigma} \cdot \hat{\boldsymbol{\theta}} U(\boldsymbol{\theta}))}{\text{Tr}(U(\boldsymbol{\theta}))} \right). \quad (\text{A10})$$

The rotationless boost that transforms a particle of mass m at rest to total momentum \mathbf{p} can be labeled by the final four velocity $\mathbf{q} := \mathbf{p}/m$:

$$A \rightarrow B(\mathbf{q}) = \sigma_0 \cosh\left(\frac{\rho}{2}\right) + \boldsymbol{\sigma} \cdot \hat{\mathbf{p}} \sinh\left(\frac{\rho}{2}\right) \quad (\text{A11})$$

where $\rho = \hat{\mathbf{p}}\rho$ is the rapidity of the Lorentz boost which is related to \mathbf{q} by

$$\cosh\left(\frac{\rho}{2}\right) = \sqrt{\frac{q^0 + 1}{2}} = \sqrt{\frac{p^0 + m}{2m}} \quad (\text{A12})$$

and

$$\sinh\left(\frac{\rho}{2}\right) = \frac{|\mathbf{q}|}{\sqrt{2(q^0 + 1)}} = \frac{|\mathbf{p}|}{\sqrt{2m(p^0 + m)}}. \quad (\text{A13})$$

$B(\mathbf{q})$ satisfies

$$B(\mathbf{q}) \begin{pmatrix} m & 0 \\ 0 & m \end{pmatrix} B^\dagger(\mathbf{q}) = \begin{pmatrix} p^0 + p^3 & p^1 - ip^2 \\ p^1 + ip^2 & p^0 - p^3 \end{pmatrix} = P \quad (\text{A14})$$

where

$$p^0 = \omega(p) = \sqrt{m^2 + \mathbf{p} \cdot \mathbf{p}}. \quad (\text{A15})$$

The rotationless boost above is called the canonical boost. The inverse transformation is obtained by reversing the sign of \mathbf{q} or \mathbf{p} .

The Wigner D functions are homogeneous polynomials of degree $2j$ in the coefficients of the $SL(2, \mathbb{C})$ matrices A_{ij} :

$$D_{\mu\nu}^j(A) := \sum_{\alpha=0}^{2j} \frac{[(j+\mu)!(j-\mu)!(j+\nu)!(j-\nu)!]^{1/2}}{(j+\mu-\alpha)!\alpha!(\alpha-\mu+\nu)!(j-\nu-\alpha)!} A_{11}^{j+\mu-\alpha} A_{12}^\alpha A_{121}^{\alpha-\nu-\mu} A_{22}^{j-\mu-\alpha} \quad (\text{A16})$$

These are representations of both $SL(2, \mathbb{C})$ and $SU(2)$. The $j = 1/2$ representation is just the matrix A .

APPENDIX B: PERMUTATION OPERATOR

Using Eq. (10) twice for the bra state ${}_1\langle k, q, \alpha |$ and the ket state $|k', q', \alpha'\rangle_2$ one gets for the matrix element of the permutation operator in our partial wave basis:

$$\begin{aligned} & {}_1\langle k, q, \alpha | P |k', q', \alpha'\rangle_1 = 2 {}_1\langle k, q, \alpha | k', q', \alpha'\rangle_2 = \\ & 2 \sum_{m_1 m_2 m_3} \sum_{\mu_2 \mu_3 \mu_s} \sum_{\mu_l \mu_\lambda \mu_I} \sum_{\mu'_2 \mu'_3 \mu'_{s'}} \sum_{\mu'_{l'} \mu'_{\lambda'} \mu'_{I'}} \\ & (\lambda \mu_\lambda \frac{1}{2}, m_1, |I, \mu_I) (j, \mu, I, \mu_I | J, M) (\frac{1}{2}, \mu_2, \frac{1}{2}, \mu_3 | s, \mu_s) (l, \mu_l, s, \mu_s | j, \mu) \\ & (\lambda', \mu_{\lambda'}, \frac{1}{2}, m_2, |I', \mu_{I'}) (j', \mu', I', \mu_{I'} | J, M) (\frac{1}{2}, \mu'_2, \frac{1}{2}, \mu'_3 | s', \mu_{s'}) (l', \mu_{l'}, s', \mu_{s'} | j', \mu') \\ & \frac{\int d\hat{\mathbf{q}} d\hat{\mathbf{q}}' \frac{1}{N(\mathbf{q}', -\mathbf{q} - \mathbf{q}')} \frac{1}{N(-\mathbf{q} - \mathbf{q}', \mathbf{q})}}{\delta(k - |\mathbf{k}(\mathbf{q}', -\mathbf{q} - \mathbf{q}')|)} \frac{\delta(k' - |\mathbf{k}(-\mathbf{q} - \mathbf{q}', \mathbf{q})|)}{\mathbf{k}^2 \mathbf{k}'^2} \\ & Y_{\mu_\lambda}^{\lambda*}(\hat{\mathbf{q}}) Y_{\mu_l}^{l*}(\hat{\mathbf{k}}(\mathbf{q}', -\mathbf{q} - \mathbf{q}')) Y_{\mu_{\lambda'}}^{\lambda'}(\hat{\mathbf{q}}') Y_{\mu_{l'}}^{l'}(\hat{\mathbf{k}}(-\mathbf{q} - \mathbf{q}', \mathbf{q})) \\ & D_{m_2 \mu_2}^{\frac{1}{2}*} [R_w(B(-\mathbf{q}/2\omega_m(\mathbf{k})), (\mathbf{k}(\mathbf{q}', -\mathbf{q} - \mathbf{q}')), \omega_m(\mathbf{k}))] \\ & D_{m_3 \mu_3}^{\frac{1}{2}*} [R_w(B(-\mathbf{q}/2\omega_m(\mathbf{k})), (-\mathbf{k}(\mathbf{q}', -\mathbf{q} - \mathbf{q}')), \omega_m(\mathbf{k}))] \\ & D_{m_3 \mu'_2}^{\frac{1}{2}} [R_w(B(-\mathbf{q}'/2\omega_m(\mathbf{k})), (\mathbf{k}(-\mathbf{q} - \mathbf{q}', \mathbf{q})), \omega_m(\mathbf{k}))] \\ & D_{m_1 \mu'_3}^{\frac{1}{2}} [R_w(B(-\mathbf{q}'/2\omega_m(\mathbf{k})), (-\mathbf{k}(-\mathbf{q} - \mathbf{q}', \mathbf{q})), \omega_m(\mathbf{k}))] \\ & {}_1 \langle (t \frac{1}{2}) T | (t' \frac{1}{2}) T \rangle_2, \end{aligned} \quad (\text{B1})$$

with

$$\begin{aligned} \mathbf{k}(\mathbf{q}', -\mathbf{q} - \mathbf{q}') &\equiv \mathbf{q}' + \frac{1}{2}\mathbf{q}(1 + y_1(q, q', x)) \\ \mathbf{k}(-\mathbf{q} - \mathbf{q}', \mathbf{q}) &\equiv -\mathbf{q} - \frac{1}{2}\mathbf{q}'(1 + y_2(q, q', x)). \end{aligned} \quad (\text{B2})$$

In Eq.(B2) occur

$$y_1(q, q', x) = \frac{E_{\mathbf{q}'} - E_{\mathbf{q}+\mathbf{q}'}}{E_{\mathbf{q}'} + E_{\mathbf{q}+\mathbf{q}'} + \sqrt{(E_{\mathbf{q}'} + E_{\mathbf{q}+\mathbf{q}'})^2 - \mathbf{q}^2}} \quad (\text{B3})$$

with $x = \hat{q} \cdot \hat{q}'$, $y_2(q, q', x) = y_1(q', q, x)$ and $E_{\mathbf{q}} \equiv \omega_m(\mathbf{q})$.

Proceeding as in [41–43] one gets the following expression for the matrix element of the permutation operator P:

$${}_1\langle k, q, \alpha | P | k', q', \alpha' \rangle_1 = \int_{-1}^1 dx \frac{\delta(k - \pi_1)}{k^2} \frac{\delta(k' - \pi_2)}{k'^2} \frac{1}{N_1(q, q', x)} \frac{1}{N_2(q, q', x)} G_{\alpha\alpha'}^{BB}(q, q', x), \quad (\text{B4})$$

with

$$\begin{aligned} \pi_1 &= \sqrt{q'^2 + \frac{1}{4}q^2(1 + y_1)^2 + qq'x(1 + y_1)} \\ \pi_2 &= \sqrt{q^2 + \frac{1}{4}q'^2(1 + y_2)^2 + qq'x(1 + y_2)} \\ N_1(q, q', x) &\equiv N(\mathbf{q}', -\mathbf{q} - \mathbf{q}') \\ N_2(q, q', x) &\equiv N(-\mathbf{q} - \mathbf{q}', \mathbf{q}), \end{aligned} \quad (\text{B5})$$

and

$$\begin{aligned} G_{\alpha\alpha'}^{BB}(qq'x) &= \frac{4\pi^{3/2}}{2J+1} (-1)^{t'} \delta_{TT'} \delta_{M_T M_{T'}} \sqrt{\hat{t}\hat{t}'} \begin{Bmatrix} 1/2 & 1/2 & t \\ 1/2 & T & t' \end{Bmatrix} \sqrt{\hat{\lambda}} \\ &\sum_{\mu_2 \mu_3} \left(\frac{1}{2} \mu_2 \frac{1}{2} \mu_3 | s \mu_2 + \mu_3 \right) \\ &\sum_{\mu'_2} \left(\sum_{m_3} D_{m_3 \mu_3}^{1/2} [R_w(B(-\mathbf{q}/2\omega_m(\mathbf{k})), (-\mathbf{k}, \omega_m(\mathbf{k}))) D_{m_3 \mu'_2}^{1/2} [R_w(B(\mathbf{q}'/2\omega_m(\mathbf{k}')), (\mathbf{k}', \omega_m(\mathbf{k}')))] \right) \\ &\sum_{\mu'_3} \left(\frac{1}{2}, \mu'_2, \frac{1}{2}, \mu'_3 | s', \mu'_2 + \mu'_3 \right) \\ &\sum_{m_1} \left(\lambda_0 \frac{1}{2}, m_1, | I, m_1 \right) D_{m_1 \mu'_3}^{1/2} [R_w(B(-\mathbf{q}'/2\omega_m(\mathbf{k}')), (-\mathbf{k}', \omega_m(\mathbf{k}')))] \\ &\sum_{\mu} (l, \mu - \mu_2 - \mu_3, s, \mu_2 + \mu_3 | j, \mu) (-)^{\mu - \mu_2 - \mu_3} Y_{l - (\mu - \mu_2 - \mu_3)}(\hat{\mathbf{p}})(j, \mu, I, m_1 | J, \mu + m_1) \\ &\sum_{\mu'} (l', \mu' - \mu'_2 - \mu'_3, s', \mu'_2 + \mu'_3 | j', \mu') Y_{l' \mu' - \mu'_2 - \mu'_3}(\hat{\mathbf{p}}')(j', \mu', I', \mu + m_1 - \mu' | J, \mu + m_1) \\ &\sum_{m_2} (\lambda', \mu + m_1 - \mu' - m_2, \frac{1}{2}, m_2 | I', \mu + m_1 - \mu') \\ &D_{m_2 \mu_2}^{1/2} [R_w(B(-\mathbf{q}/2\omega_m(\mathbf{k})), (\mathbf{k}, \omega_m(\mathbf{k}))) Y_{\lambda' \mu + m_1 - \mu' - m_2}(\hat{\mathbf{q}}'). \end{aligned} \quad (\text{B6})$$

We use standard notation $\hat{l} \equiv 2l + 1$. It is assumed that z-axis is along \mathbf{q} and the momentum \mathbf{q}' lies in the x-z plane what leads to the following components of the \mathbf{q} , \mathbf{q}' , \mathbf{k} , and \mathbf{k}' vectors

$$\begin{aligned} \mathbf{q} &= [0, 0, q] \\ \mathbf{q}' &= [q' \sqrt{1 - x^2}, 0, q'x] \\ \mathbf{k} &= [q' \sqrt{1 - x^2}, 0, q'x + \frac{1}{2}q(1 + y_1(q, q', x))] \end{aligned}$$

$$\mathbf{k}' = [-\frac{1}{2}q'(1 + y_2(q, q', x))\sqrt{1 - x^2}, 0, -q - \frac{1}{2}q'(1 + y_2(q, q', x))x] \quad (\text{B7})$$

Though the direct evaluation of Euler angles as arguments of the Wigner D-functions could be used like in [20], here we use the $SL(2, \mathbb{C})$ representations of Lorentz transformations discussed in Appendix A. This leads to

$$\begin{aligned} D^{1/2}[R_w(B(-\mathbf{q}/M_0), (\mathbf{k}, m))] &= B(\mathbf{p}_2/m)B(-\mathbf{q}/M_0)B(\mathbf{k}/m) \\ &= \sqrt{\frac{E_0 + M_0}{2M_0}} \sqrt{\frac{\omega(k) + m}{\omega(p_2) + m}} \\ &\quad - \frac{\mathbf{k} \cdot \mathbf{q}}{\sqrt{2M_0(E_0 + M_0)(\omega(k) + m)(\omega(p_2) + m)}} \\ &\quad + i\mathbf{k} \times \mathbf{q} \cdot \boldsymbol{\sigma} \frac{1}{\sqrt{2M_0(E_0 + M_0)(\omega(k) + m)(\omega(p_2) + m)}} \end{aligned} \quad (\text{B8})$$

where

$$E_0 = \sqrt{M_0^2 + q^2} \quad (\text{B9})$$

$$M_0 = 2\omega(k) \quad (\text{B10})$$

$$\mathbf{p}_2(\mathbf{k}, -\mathbf{q}) = \mathbf{k} - \mathbf{q} \left(\frac{\omega(k)}{M_0} - \mathbf{k} \cdot \mathbf{q} \frac{1}{M_0(E_0 + M_0)} \right). \quad (\text{B11})$$

-
- [1] R.B. Wiringa, V.G.J. Stoks, R. Schiavilla, Phys. Rev. C**51**, 38 (1995).
- [2] R. Machleidt, F. Sammarruca, and Y. Song, Phys. Rev. C**53**, R1483 (1996).
- [3] V.G.J. Stoks, R.A.M. Klomp, C.P.F. Terheggen, J.J. de Swart, Phys. Rev. C**49**, 2950 (1994).
- [4] J.L. Friar et al., Phys. Lett. B**311**, 4 (1993).
- [5] A. Nogga, D. Hüber, H. Kamada, and W. Glöckle, Phys. Lett. B**409**, 19 (1997).
- [6] H. Witała, W. Glöckle, D. Hüber, J. Golak, and H. Kamada, Phys. Rev. Lett. **81**, 1183 (1998).
- [7] H. Witała et al., Phys. Rev. C**63**, 024007 (2001).
- [8] K. Sekiguchi et al., Phys. Rev. C**65**, 034003 (2002).
- [9] E. Epelbaum et al., Phys. Rev. C **66**, 064001 (2002).
- [10] S.A. Coon et al., Nucl. Phys. A**317**, 242 (1979); S.A. Coon and W. Glöckle, Phys. Rev. C**23**, 1790 (1981).
- [11] B.S. Pudliner et al., Phys. Rev. C**56**, 1720 (1997).
- [12] K. Ermisch et al., Phys. Rev. C**68**, 051001(R) (2003).
- [13] K. Ermisch et al., Phys. Rev. C**71**, 064004 (2005).
- [14] Y. Maeda et al., Phys. Rev. C**76**, 014004 (2007).
- [15] W.P. Abfalterer et al., Phys. Rev. Lett. **81**, 57 (1998).
- [16] H. Witała et al., Phys. Rev. C**59**, 3035 (1999).
- [17] H. Ekstein, Phys. Rev. **117**, 1590 (1960).
- [18] W. Glöckle and W. Polyzou, Few-Body Systems, **9**, 97 (1990).
- [19] F. Coester, Helv. Phys. Acta **38**, 7 (1965).
- [20] H. Witała et al., Phys. Rev. C**71**, 054001 (2005).
- [21] H. Kamada, W. Glöckle, J. Golak, and Ch. Elster, Phys. Rev. C**66**, 044010 (2002).
- [22] W. Glöckle, T-S. H. Lee, and F. Coester, Phys. Rev. C**33**, 709 (1986).
- [23] J.L. Forest, V.R. Pandharipande, J. Carlson, and R. Schiavilla, Phys. Rev. C**52**, 576 (1995).
- [24] H. Kamada, W. Glöckle, Phys. Rev. Lett. **80**, 2547 (1998).
- [25] F. Coester, S.C. Pieper, F.J.D. Serduke, Phys. Rev. C**11**, 1 (1975).
- [26] T. W. Allen, G. L. Payne, W. N. Polyzou Phys. Rev. C**62**, 054002 (2000).
- [27] H. Witała et al., Phys. Lett. B**634**, 374 (2006).
- [28] R. Skibiński et al., Eur. Phys. J. A**30**, 369 (2006).
- [29] G.A. Miller and A. Schwenk, Phys. Rev. C**76**, 024001 (2007).
- [30] L.L. Foldy, Phys. Rev. **122**, 275 (1961); R.A. Krajcik and L.L. Foldy, Phys. Rev. D**10**, 1777 (1974).
- [31] E. P. Wigner, Ann. Math. **40**,149 (1939).
- [32] B.D. Keister and W.N. Polyzou, Adv. Nucl. Phys. **20**, 225 (1991).
- [33] P. Moussa and R. Stora in Lectures in Theoretical Physics Vol. VIIA, Ed. W. E. Brittin and A. O. Barut, Univeristy of Colorado Press, 1965.

- [34] W. Glöckle, H.Witała, D.Hüber, H.Kamada, J.Golak, Phys. Rep. **274**, 107 (1996).
- [35] W.Glöckle, The Quantum Mechanical Few-Body Problem, Springer-Verlag 1983.
- [36] P. A. M. Dirac, Rev. Mod. Phys. 21,392(1949).
- [37] T. Kato, “Perturbation theory for linear operators”, Springer, 1966, p. 543.
- [38] F. Coester and W. N. Polyzou, Phys. Rev. D26, 1348(1982).
- [39] H. Baumgärtel and M. Wollenberg, Mathematical Scattering Theory, Birkhauser, 1983 p246.
- [40] H. Kamada, H., and W. Glöckle, Phys. Lett. **B655**, 119 (2007).
- [41] R. Balian, et al., Il Nuovo Cim. **B2**, 403 (1969).
- [42] B. D. Keister and W. N. Polyzou, Phys. Rev. **C73**, 014005 (2006).
- [43] H.-C. Jean, G.L. Payne, and W. N. Polyzou, Few-Body Systems **16**, 17 (1994).
- [44] H. J. Melosh, Phys. Rev. **D9**, 1095 (1974).
- [45] P. L. Chung, F. Coester, B. Keister, and W. N. Polyzou, Phys. Rev. **C37**, 2000 (1988).
- [46] W.-M. Yao et al., Particle Data Group, J. Phys, **G 33**,1(2006).
- [47] T. Lin, Ch. Elster, W. N. Polyzou, W. Glöckle, Phys. Rev. **C76**, 014010 (2007).
- [48] H.Witała, T.Cornelius and W.Glöckle, Few-Body Syst. **3**, 123 (1988).
- [49] W. Tornow, H. Witała, Nucl. Phys. **A637**, 280 (1998).
- [50] W. Tornow, W. et al., Phys. Lett. **B 257**, 273 (1991).
- [51] J. Cub, E. Finckh, H. Friess, G. Fuchs, K. Gebhardt, K. Geissdörfer, R. Lin, and J. Strate, Few-Body Systems **6**, 151 (1989).
- [52] S.N. Bunker et al., Nucl. Phys. **A113**, 461 (1968).
- [53] H. Shimizu et al., Nucl. Phys. **A382**, 242 (1982).
- [54] E. Epelbaum, Prog. Part. Nucl. Phys. **57**, 654 (2006).