


Lecture #3: Mirror Force, Adiabatic Invariance, Polarization Drift, and Collisions

I. Magnetic Moment and the Mirror Force

- A. Magnetic Moment: 1. A current loop has magnetic moment  $\mu = IA$
2. The current loop due to Larmor motion gives



$$\mu = \frac{m v_{\perp}^2}{2 B} \quad \text{Magnetic Moment}$$

B. The Mirror Force:  $\nabla B \parallel \underline{B}$

1. Since  $\nabla \cdot \underline{B} = 0$  must always be satisfied, if the magnitude of  $B$  changes along the field line, it must involve a convergence or field lines:


- a. Consider an axisymmetric ( $\frac{\partial}{\partial \phi} = 0$ ) system in cylindrical coordinates:

$$\nabla \cdot \underline{B} = \frac{1}{r} \frac{\partial}{\partial r} (r B_r) + \frac{\partial B_z}{\partial z} = 0$$

b. Thus,  $\frac{\partial}{\partial r} (r B_r) = -r \frac{\partial B_z}{\partial z}$

- c. Let us assume  $\frac{\partial B_z}{\partial z}$  is independent of  $r$  (valid for small  $r$ ), we can integrate to yield

$$B_r = -\frac{r}{2} \frac{\partial B_z}{\partial z}$$



Thus, increasing the field magnitude along  $z$  requires a  $B_r$  component.

- e. The Lorentz force (for  $\underline{E} = 0$ ) is  $\underline{F} = q \underline{v} \times \underline{B}$ .

The  $B_r$  component of the field crossed with  $v_{\perp}$  leads to a force along the field (in the  $z$  direction).

# Lecture #3 (Continued)

Pages ②


Z. B<sub>0</sub> (Continued)

## 2. The Magnetic Mirror Force

$$F_z = -\mu \frac{\partial B_z}{\partial z}$$

z is direction along the magnetic field.

- a. Accelerates the particle along the field line in the direction of decreasing field magnitude
- b. Force confines charged particles in weak B-field regions
- c. Analogous to the electrostatic force on a charge,  $F = -q \nabla \phi$ .  
(here, magnetic field magnitude  $|B|$  behaves like a potential)




## 3. Adiabatic Invariance of Magnetic Moment

- a. As a charged particle moves through a spatially or temporally varying B field, it can be shown that magnetic moment is conserved.


$$\frac{dM}{dt} = 0$$

- b. This invariance occurs when the change in the field occurs slowly compared to fast Larmor motion. ( $\tau \gg \frac{1}{\omega_c}$ , or  $r_L \ll L$ )

## C. Confinement by a Magnetic Mirror




2. Particles are confined by mirror force



# Lecture #3


Hanes (3)

I.C. (Continued)



## 3. Pitch Angle

a. Definition: Pitch Angle,  $\alpha \equiv$  angle between velocity vector & magnetic field.



$$v_{||} = v \cos \alpha$$

$$v_{\perp} = v \sin \alpha$$

$$\underline{v} \cdot \underline{B} = vB \cos \alpha$$

4. Conservation of Energy:  $E = \frac{1}{2} m v_{||}^2 + \frac{1}{2} m v_{\perp}^2 = \text{constant}$

a. Magnetic Moment  $\mu = \frac{m v_{\perp}^2}{2B} = \text{constant}$

Function of distance along magnetic field, "s"


b. We can write  $\frac{1}{2} m v_{\perp}^2 = \mu B$ , so

$$\boxed{E = \frac{1}{2} m v_{||}^2 + \mu B}$$



constant

constant

## c. Energy Interpretation of Mirror Force



## 5. Evolution of pitch angle $\alpha$ along parallel coordinate s



a.  $E = \frac{1}{2} m v_{||}^2 + \mu B = \text{constant}$

b. As  $B(s)$  increases, conservation of energy implies  $v_{||}(s)$  decreases

c. Eventually,  $v_{||}(s) \rightarrow 0$ , and particle turns around.

# Lecture #3

Homes (4)

## I. (Continued)

### D. Loss Cone

1.  $E = \frac{1}{2} m v^2 = \text{constant}$ , so  $v = \text{constant}$ .

2. 
$$E = \underbrace{\frac{1}{2} m v^2}_{=E} \cos^2 \alpha + \mu B \Rightarrow E(1 - \cos^2 \alpha) = \mu B$$


3. Thus, 
$$\boxed{\sin^2 \alpha(s) = \frac{\mu B(s)}{E}}$$

4. Particle is mirrored when  $\alpha \rightarrow \frac{\pi}{2}$ , or  $\frac{\mu B(s)}{E} \rightarrow 1$ .

a. But, there is a maximum value to  $B(s)$ ,  $B_{\max}$ .

b. If  $E > \mu B_{\max}$ , then particle is never mirrored  $\rightarrow$  it escapes through the "neck" of the magnetic bottle.

c. But, the value of  $\mu$  depends on  $v_{\perp}$ , or the pitch angle  $\alpha$ .



a. Thus, particles with pitch angles below a threshold value will always be lost.


b. 
$$\boxed{\sin^2 \alpha_{\min} = \frac{B_{\min}}{B_{\max}}}$$

loss cone angle.

c. Particles with  $\alpha < \alpha_{\min}$  at  $B_{\min}$  will be lost!

### A. Loss Cone Distribution


Particles confined in a magnetic mirror will lead to a "loss cone distribution."



## Lecture #3 (Continued)

### I. E. Magnetosphere:

1. The Dipole Field is a natural magnetic mirror configuration



Hines ⑤

2. In the inner magnetosphere, one observes a loss cone distribution.

### II. Adiabatic Invariance:


#### A. General Concepts

1. For any periodic motion, there exists a corresponding adiabatic invariant, related to the conservation of an action integral over the periodic motion from Hamiltonian mechanics.
2. In simple magnetic field configurations (magnetic mirror, magnetic dipole), there exist a hierarchy of characteristic periodic motions, leading to a hierarchy of adiabatic invariants.
3. The invariance occurs when changes to the system occur slowly (either temporally or spatially) compared to the fast periodic motion.

#### B. Periodic Motion and Adiabatic Invariants: Magnetic Mirror


Three types of periodic motion in an axisymmetric magnetic mirror:

##### 1. Larmor Motion:




$$1^{st}: \mu = \frac{mv_{\perp}^2}{2B}$$

##### 2. Parallel Bounce Motion: (Mirror Force)



$$2^{nd}: J = m \oint v_{\parallel} ds$$

##### 3. Azimuthal Drift Motion: ( $\nabla B$ & curvature drifts)



3<sup>rd</sup>: Magnetic Flux enclosed by drift orbit remains constant

# Lecture #3 (Continued)

Howes 6

## II, (Continued)

C. What types of periodic motion exist in a dipole magnetic field? (see HW#1)

## III. Polarization Drift: Slowly varying $\underline{E}$ and constant $\underline{B}$

### A. Multiple Timescale Analysis

1. Consider a case in which the timescale of the electric field variation  $\tau \sim |\underline{E}| / |\frac{d\underline{E}}{dt}|$  is slow compared to Larmor motion,  $\tau \gg \frac{1}{\Omega}$


2. We can solve for the motion, order by order, in terms of the small expansion parameter  $\epsilon \sim \frac{1}{\tau\Omega} \ll 1$ , to obtain

$$\underline{v} = \underbrace{v_L [\cos(\Omega t) \hat{x} - \sin(\Omega t) \hat{y}]}_{\text{Zeroth-order Larmor Motion}} + \underbrace{\frac{\underline{E}(t) \times \underline{B}}{B_0^2}}_{\text{First-order } \underline{E} \times \underline{B} \text{ drift}} + \underbrace{\frac{1}{\Omega B_0} \frac{d\underline{E}}{dt}}_{\text{Second-order Polarization drift}}$$

2. Define: Polarization Drift

$$\underline{v}_p \equiv \frac{1}{\Omega B} \frac{d\underline{E}}{dt}$$

a. NOTE:  $\frac{1}{\Omega B} = \frac{m}{qB^2}$ , so polarization drift depends on charge  $\Rightarrow$  ions & electrons drift in opposite directions



4. Since electrostatic force is  $\underline{F} = q\underline{E}$ , and velocity is in direction of  $\underline{E}$ , the  $\underline{F} \cdot \underline{v} \neq 0 \Rightarrow$  polarization drift changes particle energy.

## IV. Collisions



A. 1. Another physical process that affects the motion of a single charged particle is collisions with other charged particles.

### B. Single Large-Angle vs. Many Small-Angle Collisions

1. Def: Collision time,  $T_c \equiv$  time required for particle trajectory to be deflected by  $\frac{\pi}{2}$ .

2. Deflection may be accomplished in different ways:

a. Single large-angle collision or b. Many small-angle collisions



c. Because the Coulomb force is long-range, and many particles fall within the Debye sphere and can interact with the particle, Small-angle collisions dominate over large-angle collisions!

d. It is important to remember that, within a single mean free path  $\lambda_m$  (defined as  $\lambda_m = v_f T_c$ ), the particle actually experiences many small angle collisions.

### C. Collision Frequency:

1. For a fully ionized hydrogenic (protons & electrons) plasma with density  $n_0 = n_{i0} = n_{e0}$  and temperatures  $T_i$  and  $T_e$ ,

$$\nu_{e-i} = \frac{e^4}{2^{5/2} \pi \epsilon_0^2 m_e^{1/2}} \frac{n_0}{T_e^{3/2}} \ln N_D$$

Collisions of electrons on ions

Plasma parameter,  $N_D = \frac{4\pi}{3} n_0 \lambda_D^3$

## IV. C. (Continued)

2. a. Note, because of the logarithmic, the dependence of the collision frequency on  $N_D$  is very weak
- b. Typically,  $\ln N_D \sim 10-25$  for a wide range of plasmas.

3. The important dependencies are  $\nu_{ei} \propto \frac{n_0}{T_e^{3/2}}$

a. More density plasma is more collisional.

b. Hotter plasma is less collisional  $\rightarrow$  counterintuitive!

c. Because space and astrophysical plasmas are usually very tenuous (low density) and hot, they are typically weakly collisional.

D. The "Fluid" limit of Plasmas

1. The strongly collisional limit, where  $\lambda_m$  is much smaller than all other scales of interest (typically system size  $L$  and Larmor radius  $r_L$ ), corresponds to the usual fluid limit of hydrodynamics. (more on this in Lecture #4)  $\left(\lambda_m \ll L, r_L\right)$
2. How can we possibly describe a weakly collisional plasma as a fluid?
  - a. In a magnetized plasma, the motion of particles perpendicular to  $\mathbf{B}$  is constrained by the Larmor radius  $\Rightarrow$  therefore, these perpendicular dynamics may be adequately described as a fluid.
  - b. The mean free path refers only to motions parallel to  $\mathbf{B}$ .

E. Resistivity:

1. Like-particle collisions (e-e, i-i) conserve momentum, so resistivity (resistance to current flow) is determined by electron-ion collisions  $\nu_{e-i}$ .